

Review

Stimulus-specific responses in innate immunity: Multilayered regulatory circuits

Stefanie Luecke,^{1,2} Katherine M. Sheu,^{1,2} and Alexander Hoffmann^{1,*}¹Department of Microbiology, Immunology and Molecular Genetics (MIMG), and Institute for Quantitative and Computational Biosciences (QCB), University of California, Los Angeles, Los Angeles, USA²These authors contributed equally*Correspondence: ahoffmann@ucla.edu<https://doi.org/10.1016/j.immuni.2021.08.018>

SUMMARY

Immune sentinel cells initiate immune responses to pathogens and tissue injury and are capable of producing highly stimulus-specific responses. Insight into the mechanisms underlying such specificity has come from the identification of regulatory factors and biochemical pathways, as well as the definition of signaling circuits that enable combinatorial and temporal coding of information. Here, we review the multi-layered molecular mechanisms that underlie stimulus-specific gene expression in macrophages. We categorize components of inflammatory and anti-pathogenic signaling pathways into five layers of regulatory control and discuss unifying mechanisms determining signaling characteristics at each layer. In this context, we review mechanisms that enable combinatorial and temporal encoding of information, identify recurring regulatory motifs and principles, and present strategies for integrating experimental and computational approaches toward the understanding of signaling specificity in innate immunity.

INTRODUCTION

Immune sentinel cells such as macrophages and fibroblasts initiate and orchestrate immune responses to pathogen invaders or tissue injury. As first responders, they provide finely tuned immune activation to provide pathogen-appropriate protection, while avoiding excessive tissue damage and inflammation. Recent studies have revealed that immune sentinel cells produce gene expression responses that are remarkably more stimulus- and pathogen-specific (Amit et al., 2009; Cheng et al., 2017, 2019; Sheu et al., 2019) than previously thought (Nau et al., 2002). These support the theoretical consideration that because many immune-response gene products are intrinsically toxic to the host, they should only be expressed as needed.

Immune sentinel cells respond to hundreds of ligands, including PAMPs present on or shed by pathogens, DAMPs from damaged host cells, and secreted cytokines, using dozens of pattern recognition receptors (PRRs) and numerous cytokine receptors. They reside in diverse cellular locations, such as the cell surface (e.g., Toll-like receptors [TLRs] and cytokine receptors such as TNFR, IL-1R, and IFNAR), the endosome (TLRs), and the cytosol (e.g., the nucleic acid sensing cGAS and RIG-I like receptors [RLRs]) (Boraschi et al., 2018; Dostert et al., 2019; Janeway and Medzhitov, 2002; Takeuchi and Akira, 2010; de Weerd and Nguyen, 2012). The optimal response to each threat involves a distinct mixture of dozens of functional responses that comprise cell-intrinsic pathogen defenses, cell death control, recruitment of diverse immune effector cells, tissue remodeling, and activation of systemic and adaptive immune mechanisms (Sheu et al., 2019).

The premise of this review is that integrating these mechanisms identified by molecular immunology with principles of systems biology will lead to substantial advances in our understanding of how immune sentinel responses are regulated. We specifically review the signaling mechanisms described by the innate immunology literature and relate these mechanisms to regulatory motifs commonly described by systems biology. This review therefore aims to build bridges to connect the two fields to advance our understanding of innate immune response regulation.

CONNECTING MOLECULAR IMMUNOLOGY AND SYSTEMS BIOLOGY

Molecular immunologists have generated a wealth of knowledge about the molecular signaling network, which passes information from receptors to nuclear target genes, from identifying the components, to characterizing molecular interactions, and to understanding the quantitative dose-response relationship. These have been summarized well in recent reviews (Ablasser and Hur, 2020; Barrat et al., 2019; Brignall et al., 2019; Caruso et al., 2014; Hu and Sun, 2016; Kawai and Akira, 2011; Newton and Dixit, 2012).

In parallel, systems biologists have posited that the observed stimulus specificity may be understood in terms of two main coding strategies by which information about the extra-cellular stimulus is conveyed to nuclear target genes (Figure 1A). The combinatorial code produces stimulus-specific gene expression when different stimuli activate distinct combinations of gene expression regulators that are differentially utilized by immune response genes (Buchler et al., 2003; Cheng et al., 2017; Doyle et al., 2002; Gottschalk et al., 2016; Hoffmann, 2016; Junkin et al.,






Figure 1. Principles of combinatorial and temporal coding for stimulus-specific gene expression and the layers of the stimulus-responsive network enabling it

(A) In combinatorial encoding, receptor proximal mechanisms activate stimulus-specific effector combinations. Response genes decode this through Boolean logic gate-based regulation. Stimulus-specific activity profiles of single effector modules allow temporal encoding of information, which is decoded by gene-specific mechanisms. PRR, pattern recognition receptor; Lo-pass filter, low pass filter; Hi-pass filter, high pass filter.

(B) Regulatory strategies controlling immune sentinel cell responses in the signaling network. The stimulus-responsive network consists of multiple functionally distinguishable signaling layers: (1) ligand-receptor pairs, (2) receptor-associated adaptors, (3) ubiquitin-signaling layer, (4) kinase hubs, and (5) SDTF modules and other effectors. Biochemical and cell biological mechanisms are shared between the diverse components of each signaling layer that together enable stimulus-specific coding. Ligand-receptor interactions in different subcellular locations are regulated through ligand stability, diffusion, transport, binding chaperones, and receptor transport and turnover. Adaptor proteins scaffold the assembly of larger signaling complexes. The kinetics of adaptor complex oligomerization are a key determinant of downstream signaling. Ubiquitin chains of different linkages pass on, fine-tune, and eventually downregulate signaling. They are regulated by the complex interplay of E3 ligases and DUBs. Signaling converges on three key kinase families, which are activated through scaffolded autophosphorylation or kinase cascades with toggle switch or kinase cycle mechanisms. The effector molecules activated by kinases are transcription factors inducing gene expression or post-transcriptional regulators. They are regulated through negative and positive feedback and feed-forward loops. DUBs, deubiquitinases; Ub, ubiquitin; P, phosphoryl group.

Box 1. Toolboxes of molecular immunologists and systems immunologists**Molecular immunology: experimental methods**

Molecular immunologists can interrogate all layers of the signaling network with a variety of experimental methods. In [Note S1](#), we describe a selection of commonly used methods to measure:

- Protein abundance and turnover kinetics
- Protein-protein interactions
- Protein localization
- Kinase activity
- DNA-binding activity
- Chromatin state
- RNA abundance
- Metabolic state
- Cytokine secretion

For each, we describe assay characteristics pertaining to quantitation, sensitivity, resolution, and multiplexing, and data interpretation challenges.

Perturbation studies are key to understanding a biological system. We describe a few commonly used approaches:

- Pharmacological perturbation
- Knockdown by RNA interference
- Knockout by recombination and CRISPR
- Ectopic gene expression by transfection/retroviral transduction

For each, we describe its purpose and the advantages and caveats for interpretation.

Systems immunology: regulatory motifs

Systems biologists have described regulatory motifs with characteristic dynamical behavior. These are also found in the innate immune signaling network. In [Note S1](#), we describe some with the help of simple mathematical models:

- (1) Toggle switch versus cycle switch
- (2) Multistep reaction cascades
- (3) Boolean logic gates
 - AND gates
 - OR gates
- (4) Coherent feedforward loop
- (5) Incoherent feedforward loop
 - pulse generator
 - fold-change detector
- (6) Positive feedback loop
 - bistability
 - ultrasensitivity
- (7) Negative feedback loops
 - track input duration
 - improve responsiveness
 - pulse generator
 - oscillation generator

For each, we provide a general diagram of the reactions, the ordinary differential equations that model the system, a demonstration of key behaviors that the motif allows for, and some examples of motifs in the immune response signaling network.

2016; [Scholes et al., 2017](#)). The temporal code allows stimulus specificity in gene activation through the action of a single effector, whose stimulus-specific temporal pattern of activity is decoded differentially by target genes ([Behar and Hoffmann, 2010](#); [Hoffmann, 2016](#); [Purvis and Lahav, 2013](#); [Sheu et al., 2019](#)).

Yet, despite the significant advances made in both areas, a disconnect between the molecular immunology literature and the systems biology literature has slowed progress in understanding how immune sentinel cell responses are regulated. While early systems biology studies were focused on conceptual or proof-of-principle studies and therefore employed convenient experimental systems that do not always faithfully recapitulate complex biology, recent publications demonstrate that iteration between mathematical modeling and experimental studies can provide important biological insights when applied to primary cells, whether it be in embryonic stem cells ([Heemskerck et al., 2019](#)) or during gastrulation ([Chhabra et al., 2019](#)), epithelial cell oncogenesis ([Aikin et al., 2019](#)), cancer drug responses ([Chopra et al., 2020](#)), or epigenetic control of cell death ([Vanaja et al., 2018](#)).

However, in innate immunity research, integrated experimental and mathematical modeling studies remain rare. Though several seminal studies have been carried out in related fields, including chemosensing ([Manes et al., 2015](#); [Meier-Schellersheim et al., 2006](#); [Xu et al., 2010](#)), TRAIL signaling to apoptosis ([Albeck et al., 2008a, 2008b](#); [Spencer et al., 2009](#)), T cell receptor signaling ([Chakraborty and Weiss, 2014](#); [Ganti et al., 2020](#)), and T cell signaling ([Feinerman et al., 2008a, 2008b](#); [Voisinne et al., 2015](#)), the first signaling study employing primary macrophages has only just been reported ([Adelaja et al., 2021](#)). Still, recent ad-

vances in technology such as CRISPR-mediated gene insertion ([Kundert et al., 2019](#)), single-molecule RNA *in situ* fluorescence ([Eng et al., 2019](#); [Foreman and Wollman, 2020](#); [Xia et al., 2019](#)), microfluidics ([Junkin et al., 2016](#); [Lin et al., 2019](#)), or organoid culture ([Zhang et al., 2019a](#)) allow the generation of high-resolution, quantitative datasets that enable mathematical-model-aided systems biology studies of immune processes. These studies thereby illustrate how the wealth of molecular immunology knowledge may unravel how combinatorial and temporal codes control the stimulus specificity of innate immune responses. We offer a supplementary note describing experimental perturbation methods used by molecular immunologists to dissect the components of signaling pathways and the regulatory motifs used by systems biologists to study the behavior of signaling pathways ([Box 1](#); [Note S1](#)).

LAYERS OF CONTROL IN THE STIMULUS-RESPONSIVE NETWORK

The signaling network that mediates inflammatory and anti-pathogenic gene expression consists of hundreds of proteins with diverse biochemical functions. We categorize them into five layers of regulatory control—receptors, signaling adaptors, ubiquitin-chain regulators, signaling kinases, and effectors—and briefly describe the unifying molecular mechanisms determining signaling characteristics at each layer ([Figure 1B](#)).

The first layer of control for initiating stimulus-specific responses comprises of the factors that govern ligand-receptor interactions. Ligands have distinct half-lives of bioactivity, differential diffusion,

transport, and cell uptake characteristics. Receptors are expressed at distinct levels and sorted to distinct subcellular locations via intracellular trafficking (Bagnall et al., 2018; Chen et al., 2017) including endocytosis trafficking (Leifer and Medvedev, 2016). These processes may be modulated by signals from the tissue microenvironment and by prior immune threat exposure, providing context specificity to the first control layer (Roberts et al., 2017). A variety of proteins (e.g., TIRAP and TRAM) regulate receptor localization and availability to bind ligands (Kagan et al., 2008; Leifer and Medvedev, 2016). Ligand-receptor interaction affinity may not solely be determined by structures of ligand and receptors, but also by co-factors (e.g., CD14 for LPS-TLR4 binding) that facilitate ligand-receptor interaction or receptor oligomerization (Park and Lee, 2013; Zanoni et al., 2011). Ligand binding induces di- or multimerization of PRRs and cytokine receptors, often a prerequisite for adaptor recruitment and signal transduction. For example, TLRs dimerize upon ligand binding, and RLRs multimerize into a filament on the RNA (Ablasser and Hur, 2020; Ohto and Shimizu, 2016). The cell-to-cell variability of receptor availability is an important determinant of phenotypic diversification. Systems biology modeling has proven very helpful in probing these relationships (Feinerman et al., 2008b; Gaudet et al., 2012; Martins et al., 2017).

Within the second control layer, signaling adaptors connect ligand-bound receptors with intra-cellular signaling cascades. Important adaptors in innate immune signaling include MyD88 for all TLRs except TLR3, TRIF for TLR3 and TLR4, MAVS for RLRs, and STING for cGAS. Among cytokine receptors, IL-1 signaling is also mediated by MyD88, but the large TNFR-superfamily has distinct adaptors (e.g., TRADD) (Ablasser and Hur, 2020; Boraschi et al., 2018; Chen and Jiang, 2013; Dostert et al., 2019; Luo et al., 2019). These adaptors generally lack enzymatic activity themselves but rather scaffold or nucleate the assembly of larger signaling complexes (e.g., the Myddosome). This often occurs through direct homotypic domain interactions with their receptors (e.g., CARD domains for RLR-MAVS interaction), but may also involve second messenger molecules (e.g., cGAMP for STING signaling) (Ablasser and Hur, 2020; Cadena et al., 2019; Monie et al., 2009; Wu et al., 2013). Therefore, the precise biochemical characteristics and kinetic rate constants of receptor-adaptor interactions, adaptor oligomerization, and second messenger production, half-life, and diffusion determine the signal transduction behavior of the second control layer. Interestingly, the biochemical characteristics may be further modulated by the subcellular location of the adaptor. MyD88 for example may function from both plasma membrane and endosomal compartments and may have distinct biochemical kinetic properties.

We define the third control layer by the complex network of E3 ubiquitin ligases and deubiquitinases (DUBs). These generate and degrade, respectively, K63-linked, K48-linked, and linear (M1-linked) ubiquitin chains. K63 and linear ubiquitin chains have scaffolding function to facilitate the interaction of downstream signaling kinases and substrates, while K48 ubiquitin chains induce proteasomal degradation of target proteins (Hu and Sun, 2016). Ubiquitin chains are attached either to the E3 ligases themselves (e.g., K63-linked ubiquitin on TRAF6 in MyD88 and MAVS signaling) or to upstream adaptors or bridge proteins (e.g., K63-linked chains on RIP1 kinase in TRIF and TRADD signaling) or may be unanchored (Hu and Sun, 2016; Xia et al., 2009). Regula-

tory specificity is achieved by the activation or recruitment of specific E3 ligases; the extent and duration of their activity may be modulated, and they also differ in the type of ubiquitin linkages that they generate (Hu and Sun, 2016; Parvatiyar et al., 2018). An interesting example is LUBAC, an E3 ligase that synthesizes linear ubiquitin chains, which contribute to activation of the I κ B kinase (IKK) in response to TNF signaling but negatively affect RIG-I signaling (Hu and Sun, 2016; Tokunaga, 2013). Similarly, DUBs may have distinct roles in different pathways. CYLD, for example, attenuates RIG-I signaling by removing K63-linked chains but promotes STING signaling by removing K48-linked chains (Friedman et al., 2008; Zhang et al., 2018). Interestingly, the anti-inflammatory protein TNFAIP3/A20 was reported to act as both a DUB and an E3 ligase, negatively regulating NF- κ B signaling by removing K63-linked chains and adding K48-linked chains to multiple targets (Catrysse et al., 2014; Wertz et al., 2004), though their physiological roles have been questioned (De et al., 2014; Skaug et al., 2011). Thus, the ubiquitin control layer regulates interactions, activity, and degradation of key signaling proteins and provides many levers for fine-tuning of responses, cross-regulation, and network memory.

The fourth control layer involves three central families of response kinases, namely MAPKs (JNK, p38, ERK), the IKK complex (consisting of IKK α , IKK β , and IKK γ /NEMO), and TBK1 (Ablasser and Chen, 2019; Newton and Dixit, 2012). These kinases serve as the regulatory network hubs that directly or indirectly activate downstream effectors. Kinase activity in immune response pathways is generally initiated by trans-autophosphorylation when at least two kinase molecules are brought into proximity through oligomerized adaptors (e.g., activation of TBK1 downstream of STING) or ubiquitin chains (e.g., activation of TAK1 and IKK) (Emmerich et al., 2013; Häcker and Karin, 2006; Hu and Sun, 2016; Kensche et al., 2012; Ma et al., 2012; Polley et al., 2013; Shu et al., 2013; Xia et al., 2009; Zhang et al., 2019b)—in contrast, tyrosine kinase receptors are activated via ligand-induced conformational changes of receptor dimers (Livnah et al., 1998). Kinase activity may then be amplified through kinase cascades, the most prominent being MAPK cascades (Kholodenko and Birtwistle, 2009; Witzel et al., 2012). Scaffolds may also connect kinases to their substrates. For example, for IKK, NEMO, which connects the catalytic components to ubiquitin chains, also contains an I κ B α recruitment domain that ensures efficient activation of NF- κ B (Schröfelbauer et al., 2012). Kinase activity is then terminated via one of two mechanisms. In the classic toggle-switch model, inactivation occurs via dephosphorylation by phosphatases (Bardwell, 2008; Ferrell and Bhatt, 1997; Ferrell and Ha, 2014; Nolen et al., 2004; Zhao et al., 2012)—phosphatases may also target signaling molecules further upstream, such as signaling adaptors STING, TRIF, and MAVS (Ni et al., 2020; Xiang et al., 2016). However, IKK is thought to be regulated via a multi-state reaction cycle, in which an autophosphorylated inactive state must be recycled to a phosphorylation-free poised state before reactivation can occur (Behar and Hoffmann, 2013). The two inactivation modalities impart fundamentally different dynamic control to the kinase activity, and phosphatases play distinct roles. Their potential for inducible expression provides another level of regulatory control.

The fifth control layer comprises the signaling pathway effectors that directly control gene expression. There are two main

categories of effector proteins: transcription factors and proteins acting post-transcriptionally. Three prominent signal-dependent transcription factor (SDTF) families may be directly mapped to upstream kinases: MAPK-AP1/ATF, IKK-NF- κ B, and TBK1-IRF (Cargnello and Roux, 2011; Hayden and Ghosh, 2004; Honda et al., 2006). In addition, NFAT is activated in dendritic cells in a Ca²⁺-dependent manner (Fric et al., 2012; Zanoni and Granucci, 2012). Some kinases regulate post-transcriptional regulatory mechanisms. For example, IKK phosphorylates the mRNA decapping activator ECD4, affecting the levels of hundreds of mRNAs (Mikuda et al., 2018). Further, production of TNF involves not only NF- κ B-dependent transcription, but MAPK p38/ERK-dependent mRNA splicing, mRNA stabilization via tristetraprolin (TTP) (Mahtani et al., 2001), and pro-protein processing via TACE (Caldwell et al., 2014), which explains why TNF secretion by single cells does not correlate well with NF- κ B activity (Junkin et al., 2016). Effectors may be regulated in a simple toggle switch fashion with phosphatases terminating effector activity when kinase activity has diminished (e.g., IRFs and the MAPK substrates AP1 or TTP) (Gu et al., 2014; Long et al., 2014). Effectors may also be subject to regulatory feedback. NF- κ B is subject to complex negative feedback regulation via I κ B proteins, which inhibit NF- κ B through direct binding in the nucleus and escort NF- κ B back into the cytoplasm. This inhibition is reversible for some I κ B isoforms (I κ B α , I κ B ϵ) as their degradation is induced by IKK (Mitchell et al., 2016). In contrast, IRF induces the interferon- β (IFN- β) feedforward system to activate ISGF3-driven gene expression programs in both primary response and bystander cells (Ourthiague et al., 2015). In addition, MAPK phosphatases such as dual-specificity phosphatases (DUSPs) are strongly upregulated by MAPK-activating PRRs, contributing to negative feedback on MAPK pathways (Brondello et al., 1999; Kuwano et al., 2008; Salojin et al., 2006). The strength and time delays in these feedback systems are key to imparting specific activation dynamics to the effectors. Further quantitative studies are needed to enable reliable mathematical modeling studies.

It is the stepwise transmission and modulation of the signal by these biochemical and cell biological mechanisms through the network layers that mediates combinatorial and temporal coding of response specificity. While numerous molecular mechanisms have been described for key TLR pathways, how they function together to generate stimulus-specific responses is generally less well understood. As will be discussed below, stimuli activate combinations of effectors through the molecular mechanisms within receptor, adaptor, ubiquitin, and kinase layers, while stimulus-specific dynamic control is mediated by molecular mechanisms in all five control layers.

MECHANISMS THAT ENABLE COMBINATORIAL ENCODING

When stimuli activate different combinations of signaling pathways, they convey stimulus-information by combinatorial coding (Figure 1A). Hence, branching of the signal into two (or more) signaling pathways is key to combinatorial coding. Here, we first present the molecular mechanisms that allow for pathway branching and then describe how these may be deployed stimulus-specifically.

Pathway branching occurs at multiple control layers of the signaling network. Pathways usually branch when a regulator sequentially (not coincidentally) engages two signaling complexes, often by moving from one sub-cellular compartment to another. A well-studied example of this in the receptor layer is how TLR4 engages two different adaptors at different sub-cellular locales, MyD88 at the cell surface (activating NF- κ B and MAPKs) and TRIF in the endosomes (additionally activating IRF3) (Figure 2A). This adaptor switch is mediated by a switch of the bridge proteins TIRAP and TRAM, which facilitate the association of the adaptors with TLR4 within the plasma or endosomal membranes, respectively (Balka and De Nardo, 2019; Deguine and Barton, 2014). Thus, a key principle of combinatorial encoding by TLR4 is that adaptor engagement is sequential and may be regulated by regulating the transport of TLR4 from one compartment to the other (Leifer and Medvedev, 2016).

Signaling pathways may also branch at the adaptor layer (Figure 2B): TRIF-mediated activation of the TBK1-IRF3 axis and the IKK-NF- κ B axis is a prominent example (Chen and Jiang, 2013; Jiang et al., 2004), with MAVS and STING functioning similarly (Abe and Barber, 2014; Dunphy et al., 2018; Fang et al., 2017a, 2017b; Liu et al., 2013; de Oliveira Mann et al., 2019; Yoboua et al., 2010; Zhang et al., 2019b). On the one hand, TRIF oligomerization leads to the recruitment of RIP1 kinase, which is ubiquitinated by E3 ligases (Cusson-Hermance et al., 2005). Those ubiquitin chains lead to IKK recruitment and trans-autophosphorylation, allowing IKK to activate NF- κ B (Hu and Sun, 2016; Kensche et al., 2012; Polley et al., 2013). On the other hand, oligomerized TRIF also recruits TBK1, though the exact mechanism is not fully elucidated (Funami et al., 2008). Although initial TBK1 activation may require a priming phosphorylation through IKK activity (Abe et al., 2020), trans-autophosphorylation of TBK1 leads to its full activation (Ma et al., 2012; Shu et al., 2013). Activated TBK1 phosphorylates the adaptor, allowing IRF3 binding to the adaptor, which subsequently serves as a scaffold for TBK1-mediated IRF3 phosphorylation (Abe et al., 2020; Liu et al., 2015b). Thus, activation of two pathways by a single adaptor appears to occur sequentially, in which the second pathway activation requires a post-translational modification of the adaptor. Similarly, NF- κ B and IRF3 activation seem spatially regulated downstream of STING, with NF- κ B activation occurring from the endoplasmic reticulum (ER; in which STING is first activated), while IRF3 signaling requires STING trafficking to the Golgi (Ni et al., 2017; Stempel et al., 2019).

Ubiquitin chains constitute remarkably efficient signaling scaffolds that mediate pathway branching to the MAPK and NF- κ B pathways (Figure 2C). For example, ubiquitin chains attached to adaptor TRAF6 provide scaffolds for the trans-autoactivation of kinases, such as TAK1, a MAPK kinase kinase (MAPKKK) (Ishitani et al., 2003; Kanayama et al., 2004; Scholz et al., 2010; Schröfelbauer and Hoffmann, 2011; Xia et al., 2009). In addition, the IKK complex trans-autophosphorylates when clustered on (preferentially linear) ubiquitin chains through its ubiquitin-binding subunit NEMO (Ea et al., 2006; Haas et al., 2009; Kensche et al., 2012; Polley et al., 2013; Rahighi et al., 2009; Tokunaga, 2013; Xia et al., 2009). Thus, the ubiquitin layer provides numerous opportunities for signaling crosstalk and pathway branching.




Figure 2. Molecular mechanisms for combinatorial encoding to achieve stimulus-specific responses

(A–D) Pathway branching, in which one upstream protein activates multiple downstream signaling mediators, can occur at several signaling layers. Examples include branching (A) at the level of receptors with TLR4 activating the MyD88 and TRIF pathway; (B) at the level of adaptors with TRIF, MAVS, and STING being capable of IRF3 and NF-κB activation; (C) at the level of ubiquitin chains, which scaffold activation of TAK1 for MAPK activation and of IKK for NF-κB activation; and (D) at the level of kinase cascades which branch to activate IKK and the MAPKs p38, JNK, and ERK. (E) Combinatorial coding in the innate immune network is mediated by receptor-specific utilization of adaptors and pathway branching at the adaptor layer.

(Shim et al., 2005). Interestingly, IKK also engages in pathway branching by not only activating the IκB-NF-κB axis, but also the p105-TPL2-ERK axis (Beinke et al., 2004; Gantke et al., 2012). IKK mediated p105 degradation is essential for Tpl2 activation, which is the MAPKKK for ERK (Beinke et al., 2004; Gantke et al., 2012; Waterfield et al., 2003). Indeed, IKK may further phosphorylate Tpl2 (Roget et al., 2012). IKK’s substrate specificity seems to be controlled by NEMO, which is essential for IKK’s activation by inflammatory stimuli: NEMO directs IKK’s substrate specificity to IκBα, but may then dissociate from the IKK dimer, broadening IKK’s specificity to include p105 (Schröfelbauer and Hoffmann, 2011; Schröfelbauer et al., 2012). Furthermore, in pDCs, IKK was reported to activate IRF7 downstream of MyD88 mediating TLR7 and TLR9 signals (Honda et al., 2004; Hoshino et al., 2006; Kawai et al., 2004; Pauls et al., 2012; Uematsu et al., 2005). Spatial segregation between distinct endolysosomal compartments may contribute to the switch from NF-κB to IRF7 signaling (Honda et al., 2005; Sasai et al., 2010). This illustrates that pathway branching at the ki-

nase control level also involves sequential mechanisms and may be cell-type specific.

To mediate stimulus-specific responses, pathway branching must be deployed stimulus specifically. The combination of the IKK-NF-κB and JNK-AP1 axes is activated by all adaptor proteins discussed here (i.e., MyD88, TRIF, STING, MAVS, and TRADD) ensured by pathway branching at the ubiquitin chain and the kinase layers, and thus generates a common core gene expression response (Amit et al., 2009; Cheng et al., 2017). However, originating at the receptor layer,

Given the enzymatic activity of kinases, they may phosphorylate several substrates and thus mediate pathway branching (Figure 2D): TAK1 mediates pathway branching to the MKK4/7-JNK axis and the IKK-NF-κB axis. TAK1 is the initiator of the MAPK cascade that leads to JNK activation (Cargnello and Roux, 2011). In addition, TAK1 may initiate IKK activation (Cohen and Strickson, 2017; Emmerich et al., 2013; Shim et al., 2005; Wang et al., 2001; Zhang et al., 2014), though this is stimulus-specific and may be cell-type dependent. It is not clear what determines TAK1’s substrate specificity

stimulus-dose and exposure route modulate JNK and NF- κ B signaling to allow distinction of bacterial species, dose, and infection versus bystander status (Lane et al., 2019). More obviously, the TBK1-IRF3 and the MAPKp38 axes are deployed stimulus specifically by virtue of quantitative differences because of receptor-specific utilization of distinct adaptors, followed by pathway branching at the adaptor layer. Their stimulus-specific activation in conjunction with the common core pathways thus enables combinatorial coding (Figure 2E).

The TBK1-IRF3 axis mediates stimulus-specific induction of type I IFN (Kawai and Akira, 2011). In macrophages (and conventional dendritic cells), the IRF pathway is activated only in response to endosomal signaling by TLR4 and TLR3, and cytosolic signaling from cGAS and RLRs (Caruso et al., 2014; Kawai and Akira, 2011; Sparrer and Gack, 2015). This specificity is mediated by the ability of TLR3 and TLR4 to engage the TRIF adaptor in the endosome, while cGAS and RLRs engage STING and MAVS, respectively. TRIF, STING, and MAVS share an activation mechanism for the TBK1-IRF3 axis, while also being able to activate IKK-NF- κ B and TAK1-JNK axes. Thus, for example, early studies showed TLR4 activation by lipopolysaccharide (LPS) from Gram-negative bacteria activates both NF- κ B and IRF3, while TLR2 activation by peptidoglycans from Gram-positive bacteria lacks the IRF3 component (Doyle et al., 2002). As IRF3 induces expression of the cytokine IFN- β , which activates the ISGF3 TF complex in an autocrine and paracrine manner, the effect of stimulus-specific activation of IRF3 is dramatically amplified both within the first-responder cell and its neighbors.

A second, less well-characterized example is the stimulus-specific activation of MAPKs p38 and ERK. Several lines of evidence indicate that functionally potent MAPKp38 activation is stimulus specific, even if some level of activity can be detected in response to many stimuli, such as TLR ligands, cytosolic nucleic acids, and TNF. First, transcriptome profiling of fibroblasts and macrophages stimulated with a diverse set of PAMP and cytokine ligands revealed a set of about 80 genes that required both NF- κ B and MAPKp38 activity (Cheng et al., 2017). Second, MAPKs p38 and ERK have a key role in post-transcriptional processing, mRNA half-life, pro-protein processing, and secretion of TNF—cell stimulation with TNF failed to activate these processes (Caldwell et al., 2014). Finally, while stimulation with high concentrations of LPS led to TNF production, low concentrations activated NF- κ B but failed to produce TNF (Gottschalk et al., 2016), in line with the conclusion that single-cell TNF secretion was better correlated with MAPK activation than NF- κ B (Junkin et al., 2016). However, the mechanism of stimulus-specific activation of the MAPKp38 axis remains poorly understood. One hypothesis is that the MyD88 oligomerization into the Myddosome complex results in a thresholded dose response (Cheng et al., 2015) such that p38 is only activated at high ligand concentrations. In addition, the MAPK cascade itself is thought to have a thresholded dose response curve (Huang and Ferrell, 1996), dependent on the level of phosphatase activity (Altan-Bonnet and Germain, 2005). Interestingly, MAPKp38 activity may be induced through two different pathways, namely MKK4 downstream of TAK1 MAPKKK activity and MKK3/6 activated by Tpl-2 (Cohen and Strickson, 2017; Pattison et al., 2016). This may suggest that while TNF induces MAPKp38 via MKK3/6,

LPS engages both pathways to generate a more potent p38 activity (Pattison et al., 2016).

In sum, combinatorial effector activation may encode information about both the molecular identity and the dose of a stimulus. Quantitative differences in the dose response curves of co-activated pathways increase the capacity for producing stimulus-specific responses via combinatorial coding. Future studies should address not only the mechanisms of pathway branching (e.g., the role of IKK in TBK1-IRF activation and the mechanisms of NF- κ B and MAPK activation downstream of cytosolic nucleic acid sensing remain unclear), but also address dose-response behavior of each of the branched pathways and whether they are modulated by the microenvironment and the cell's exposure history. Importantly, during the infection process, pathogens stimulate several PRRs and auto-crine cytokines (which further activate related pathways) that, in aggregate, characterize pathogen-specific responses *in vivo* (Gottschalk et al., 2019). Given that *in vitro* studies with multiple ligands revealed instances of synergy or antagonism between pathways (Gottschalk et al., 2019; Gutschow et al., 2019; Lin et al., 2017; Pandey et al., 2020; Tan et al., 2014), further study of cross-regulatory connections between co-activated pathways is warranted.

MECHANISMS THAT ENABLE TEMPORAL ENCODING

Signaling involves the transient perturbation of a homeostatic steady state. Hence, the activity of a signaling effector exhibits a temporal trajectory. The observation that two stimuli may trigger distinct temporal trajectories, and thereby differential gene expression, led to the hypothesis of a “temporal signaling code” (Figure 1A) (Hoffmann and Baltimore, 2006). Temporal trajectories may be quantitatively characterized by a set of features, such as peak amplitude, fold change, amplitude of the second signaling phase, total duration, or oscillations (Behar and Hoffmann, 2010; Levine et al., 2013; Li and Elowitz, 2019; Purvis and Lahav, 2013). Here, we describe how such dynamic features may be generated, or encoded, by the successive modulation of signaling activity by multiple mechanisms as the signal passes through successive control layers.

Within the first control layer, the first determinant of signaling dynamics is ligand availability as determined by ligand diffusion, endocytosis, and decay. Secreted TNF, for example, has a short half-life, whereas the endotoxin LPS persists. In tissue environments, diffusion rates and rates of uptake by other cells may also be important (Bagnall et al., 2018). Next, receptor clustering upon ligand binding (Caré and Soula, 2011) may threshold the dose-response curve preventing spurious activation in the absence of ligand (Bray et al., 1998). Following ligand binding, endocytosis of receptor-ligand complexes reduces the availability of receptors for further stimulation. The speed of receptor replenishment, through recycling or *de novo* synthesis, together with ligand half-life is a key determinant of the amplitude of second phase signaling (Becker et al., 2010) (Figure 3A).

In the next layer, the kinetic characteristics of adaptors are important determinants of signaling dynamics. For example, stepwise assembly of the Myddosome imparts the MyD88 pathway a thresholded, close-to-digital dose-response curve,




Figure 3. Molecular mechanisms for temporal encoding to achieve stimulus-specific responses

(A–D) Mechanisms contributing to important features of temporal signaling dynamics: (A) the second phase amplitude is controlled by ligand stability, receptor turnover dynamics, and kinase activity kinetics. (B) Dose-to-peak-amplitude-encoding is dependent on adaptor oligomerization properties, which differ between MyD88 and TRIF. (C) Oscillations in effector activity trajectories of e.g., NF- κ B or MAPKp38 are introduced by negative feedback loops with delay. (D) Total duration of signaling is modulated by positive feedback or irreversible negative feedback, such as the induction of RelA and I κ B δ expression, respectively.

of TNF signaling and therefore limits the weaker signals of later time points. Both induced A20 feedback expression and constitutive A20 expression at certain levels can fulfill this function (Werner et al., 2008).

The kinase layer provides for interesting temporal modulation of signaling activity. The two-state toggle switch kinase transmits the incoming signal, but there is an inherent trade-off between faithful turn-off with cessation of the signal (requiring high phosphatase activity) and reduction of the peak amplitude dynamic range. This is resolved by kinase cascades (e.g., in the MAPK pathway), which amplify the signal's dynamic range without losing too much temporal control (Heinrich et al., 2002; Kholodenko and Birtwistle, 2009). An alternative topology is the kinase cycle (e.g., canonical IKK). This allows for a strong peak of activation, which triggers the kinase's transition to an inactive state (Delhase et al., 1999). From this, the kinase has to be recycled (via a phosphatase) to a poised state

before reactivation. This results in a two-phased temporal trajectory with the duration of the first phase determined by the inactivation reaction and the amplitude of the second phase determined by the recycling reaction (Behar and Hoffmann, 2013). Thus, the distinction between first versus second signaling phases is enforced by kinase cycles or diminished by toggle kinase cascades (Behar and Hoffmann, 2013) (Figure 3A). When positive feedback is included in the kinase layer, an excitable dose-response curve may produce digital all-or-none responses (Shinohara et al., 2014).

Within the ubiquitin layer, complex chain regulatory dynamics and their interplay with ubiquitin-binding proteins likely also control signaling; however, a quantitative, time-resolved understanding requires further study. Anti-inflammatory A20, one prominent regulator of ubiquitin-chain dynamics, blunts the dose response

whereas TRIF dimerization leads to a more linear dose-response curve and translates ligand concentration into effector amplitude more faithfully (Cheng et al., 2015) (Figure 3B). Interestingly, Myddosome-mediated signaling is transient and limited to a few hours, but how it is turned off remains unclear. In the case of IL-1 signaling, kinase mediator IRAK4 is proteolyzed after signaling (Ferrao et al., 2014), which may limit signal duration, but how Myd88 signaling is temporally limited demands further study. In the case of TRIF-signaling, the progressive acidification of the endosome appears to be a determinant of its duration, allowing for long-lasting and variable signal trajectories (Cheng et al., 2015; Husebye et al., 2006; McGettrick and O'Neill, 2010).

Within the ubiquitin layer, complex chain regulatory dynamics and their interplay with ubiquitin-binding proteins likely also control signaling; however, a quantitative, time-resolved understanding requires further study. Anti-inflammatory A20, one prominent regulator of ubiquitin-chain dynamics, blunts the dose response

effector activity if the input signal to the feedback loop persists (Hoffmann et al., 2002; Nelson et al., 2004; Tomida et al., 2015) (Figure 3C). The IKK- $\text{I}\kappa\text{B}\alpha$ -NF- κB system generates particularly robust oscillations because of the rapid NF- κB -responsive $\text{I}\kappa\text{B}\alpha$ synthesis (including an appropriate delay to the negative feedback), rapid nuclear import of $\text{I}\kappa\text{B}\alpha$, efficient removal NF- κB from the DNA through $\text{I}\kappa\text{B}\alpha$ binding, rapid nuclear export of the complex, and responsiveness to IKK, which allows for efficient $\text{I}\kappa\text{B}\alpha$ degradation after each cycle of re-synthesis (Mitchell et al., 2016). A delay in the negative feedback (e.g., $\text{I}\kappa\text{B}\epsilon$) can diminish oscillatory propensity (Kearns et al., 2006). Irreversible negative feedback, as is mediated by A20 or by $\text{I}\kappa\text{B}\delta$ (which, unlike other $\text{I}\kappa\text{B}$ isoforms, is not targeted for degradation by canonical IKK) can lead to a shutdown or attenuation of signaling that may impact both the amplitude of the second signaling phase, the total signal duration, and the response to a secondary signal (Shih et al., 2009; Werner et al., 2008) (Figure 3D). Hence, these proteins also mediate signaling crosstalk (Werner et al., 2008), as do the SOCS proteins regulating IRF/ISGF3 responses (Krebs and Hilton, 2001; Liu et al., 2015a).

Positive feedback and feedforward mechanisms mediated by (1) expression of signaling substrates or (2) induction of cytokines may extend the duration of effector activity. NF- κB -responsive expression of the main NF- κB dimer component, RelA, sustains long-term NF- κB activity (Sung et al., 2014) (Figure 3D), while ISGF3-inducible expression of its constituents (STAT1/2/IRF9) renders cells more responsive to subsequent IFN stimulation (Ivashkiv and Donlin, 2014). Importantly, induced cytokines not only amplify or prolong signaling in primary response cell, but also trigger signaling in previously un-exposed or un-responsive neighboring cells. For example, TNF was found to have an autocrine role in prolonging NF- κB signaling in response to synthetic TLR9 ligand CpG but primarily paracrine functions in response to LPS (Caldwell et al., 2014). Similarly, IRF3-dependent IFN- β induces amplification of type I IFN signaling in the primary response cell, but also initiates antiviral responses in neighboring cells. These cytokines are primarily controlled by a feedforward mechanism rather than positive feedback, as potent paracrine signaling coupled to positive feedback can lead to runaway inflammation (Ourthiague et al., 2015): In the case of TNF, MAPKp38 is critical for its expression, but is not strongly activated by it (Caldwell et al., 2014). In the case of IFN- β , IRF3 is critical for its expression but is not activated by it (Honda et al., 2006). That regulatory feature also allows these cytokines to extend the duration of signaling, but not indefinitely.

In order to mediate stimulus-specific responses, temporal trajectories must be stimulus specific. Thus far, this has been most extensively described for NF- κB , where the temporal trajectory may encode stimulus duration, dose, and molecular identity. Early studies examined how the duration of the TNF stimulus is encoded in the temporal profile of NF- κB . Persistent TNF stimulation leads to characteristic oscillatory NF- κB activity, while short TNF pulses lead to only a single peak whose duration of approximately 45 min is independent of stimulus pulse duration (as short as 1 min) (Hoffmann et al., 2002; Werner et al., 2008). Two temporal amplification steps explain this. First, the receptor-ligand complex and mechanisms in the ubiquitin layer

provide for at least 10 min of IKK activity, and second, the $\text{I}\kappa\text{B}\alpha$ feedback loop ensures 45 min of activity. When TNF pulses exceed 60 min, subsequent cycles of oscillations are activated allowing for faithful stimulus-duration to response-duration encoding (Hoffmann et al., 2002).

Stimulus dose can also be encoded in the temporal profile of the effector. For TNF, dose is encoded in amplitude and duration (Cheong et al., 2006, 2011; Tay et al., 2010; Turner et al., 2010). Amplitude provides only partial dose information as single-cell studies observed some degree of ultrasensitivity (Tay et al., 2010), and amplitude measurements at 30 min for doses over 3 orders of magnitude contained less than 1 bit of information (Cheong et al., 2011). Instead, the short half-life of TNF mediates dose-dependent control of NF- κB duration (Adelaja et al., 2021). For PAMPs, the MyD88 and TRIF pathways encode dose information differently (Cheng et al., 2015). MyD88-dependent PAMPs show an all-or-none response, with the threshold determined by receptor and MyD88 abundance. PAMPs engaging the TRIF pathway show a more graded dose-response with dose-dependent amplitudes. However, in both pathways the speed of response increases with dose.

The important question of whether molecular identity is represented in the temporal pattern of NF- κB activity had, until recently, received confusing answers. Early population level biochemical studies in primary fibroblasts showed that, in contrast to the oscillatory activity induced by TNF, the NF- κB response to LPS-TLR4 signaling is persistent and non-oscillatory (Covert et al., 2005; Werner et al., 2005). However, studies with immortalized cell lines did not provide confirmation at the single-cell level because NF- κB responses to several ligands showed noisy, largely oscillatory trajectories (Hughes et al., 2015). A recent study with primary macrophages provided clearer answers and identified six features within NF- κB temporal trajectories correlated with the molecular identity of the ligand (Adelaja et al., 2021). For example, whereas TNF, TLR2/1 ligand Pam3CSK4, and LPS activate NF- κB rapidly, the response to CpG and TLR3 ligand poly(I:C) was slower, due to a delay in endosomal ligand availability. Remarkably, oscillatory content was substantially higher in response to cytokine TNF than to PAMPs, largely because of the distinct amplitude of the second phase of IKK activity that either overrides $\text{I}\kappa\text{B}\alpha$ negative feedback (in case of PAMPs) or allows it to generate oscillations (in case of TNF). Each ligand generated a unique combination of these six features of the temporal NF- κB code that may allow target genes to be expressed stimulus specifically.

Thus, the dynamic features in the temporal profile of an effector, modulated by biochemical and cell biological mechanisms at every signaling layer, can encode information about the stimulus such as molecular identity, dose, and duration. While two decades of research into temporal encoding in the NF- κB pathway have provided some mechanistic explanations for a handful of ligand receptor combinations, the temporal encoding of other effectors, such as the IRF and MAPKp38, remains underexplored. Indeed, how sequentially administered pulses of stimulation or multiple simultaneously administered stimuli are registered by the temporal activity profiles of effectors can provide insights about the intrinsic dynamic regulation of the signaling pathway (Ashall et al., 2009; Gutschow et al., 2019; Kellogg and Tay, 2015).

DECODING MECHANISMS THAT ENABLE STIMULUS-SPECIFIC GENE EXPRESSION

For immune sentinels to generate stimulus-specific gene expression, the gene regulatory network must interpret the combinatorial and temporal features of effector proteins. The gene regulatory network associated with each immune response gene involves DNA-associated mechanisms that move nucleosomes, establish enhancers, catalyze chromatin looping, or recruit components of the pre-initiation complex; RNA-associated mechanisms involving transcription initiation and RNA polymerase promoter clearance, transcription elongation, mRNA processing, splicing and export, or cytoplasmic decay; and translation-associated mechanisms such as pre-protein processing or activation.

Combinatorial decoding mediates stimulus-specific gene expression when immune response genes are differentially responsive to the various signaling effectors. Transcriptome profiling revealed that genes may contain AP1 binding sites, κ B elements or ISRE sites, or combinations of these (Amit et al., 2009; Cheng et al., 2017). However, two binding sites being present does not indicate whether the gene is responsive to either pathway, or only when both pathways are active. The promiscuity of the former and specificity of the latter are aptly described by the conceptual framework of Boolean OR and AND logic gates, respectively (Buchler et al., 2003; Morris et al., 2010). Beyond combinatorial activation of signaling pathways, the number of combinatorial possibilities increases drastically when differential TF-promoter binding affinities and activation strengths are also considered (Brignall et al., 2019).

Within the pathogen-responsive transcriptome, a cluster of about 100 genes were identified as being activated by either NF- κ B or ISGF3, their regulatory regions tend to contain both κ B sites and ISREs and therefore classified as OR gate genes (Cheng et al., 2017). However, NF- κ B and ISGF3 do not have redundant roles because they show distinct temporal control: only their combined action ensures that such genes are both rapidly induced (first by NF- κ B) and expressed for extended times (then by ISGF3) (Figure 4A). Indeed, when such genes encode mRNAs with a long half-life, transient TF activity may be insufficient for full gene expression (Hao and Baltimore, 2009; Sen et al., 2020). In this regard, the OR gate may be thought of as providing an additional mechanism for temporal coding because it allows for the decoding of the combined temporal trajectory of both NF- κ B and ISGF3.

In the established conceptual framework of Boolean logic gates, AND gate logic provides for high stimulus-selectivity, as both signaling pathway effectors must be active. It was thought that synergy between two TFs is achieved either by cooperative binding of DNA or by the cooperative recruitment of co-activators such as CBP/p300, which contains multiple TF interaction domains (Merika et al., 1998). However, in the pathogen-responsive transcriptome, there is little evidence for TFs needing to present at the same time or “coincident AND gates” at gene promoters. Rather, the sequential actions of signaling pathway effectors are required (Cheng et al., 2017). The discovery of sequential Boolean gates emphasizes the importance of integrating combinatorial and temporal signaling concepts for a full understanding of decoding.

One example of sequential AND gates are the small number of genes whose transcriptional initiation is regulated by both IRF3 and NF- κ B (Freaney et al., 2013; Tong et al., 2016). While *Irf3* is the most well-known example, the molecular mechanism enabling NF- κ B and IRF3 synergy is not clearly understood (Panne et al., 2007). In the case of *Ccl5*, recent studies suggest that IRF3 may facilitate a chromatin opening step that is required for the recruitment of NF- κ B, which then activates transcription (Figure 4A) (Ramirez-Carrozzi et al., 2009; Tong et al., 2016). A second example of sequential AND gates pertains to a number of important cytokine genes, such as *Tnf*, *Il6*, and *Il1b*, that are dependent on both NF- κ B and MAPKp38 activation downstream of the MyD88 adaptor (Cheng et al., 2017). While NF- κ B is sufficient for transcriptional initiation, it appears that several post-transcriptional control steps, particularly mRNA stabilization, are p38-dependent (Figure 4A). In the case of TNF, the combined action of MAPKp38 and ERK enhance pre-mRNA splicing, cytoplasmic mRNA half-life, translation, and pro-protein processing (Caldwell et al., 2014). Thus, the sequential action of diverse biochemical processes in several different cellular compartments form a sequential AND gate in the production of secreted TNF. AND gate function may also control epigenomic reprogramming: a study on IFN- γ -induced gene repression showed that dissociation of active enhancers was a result of an AND gate at MAF-bound enhancers by IFN- γ -induced repression of MAF transcription and coordinate dissociation of PU.1 and C/EBP (Kang et al., 2017).

While combinatorial decoding interprets the activities of multiple TF pathways, temporal decoding distinguishes between distinct temporal trajectories of a single signaling effector and requires mechanisms that are kinetically regulated. For example, a kinetically slow mechanistic step not only transduces the instantaneous activity but also stores information about past activity (Behar and Hoffmann, 2010; Purvis and Lahav, 2013). Three molecular mechanisms have been identified: mechanisms controlling mRNA half-life, nucleosome displacement, and enhancer formation.

Long mRNA half-life has been shown by both mathematical modeling and experimental data to decode the duration of signaling effector activity. As short half-life transcripts achieve their half-maximal induction levels faster than long half-life transcripts, only long duration signals may allow full induction of long half-life mRNAs (Behar and Hoffmann, 2010; Hao and Baltimore, 2009). In the pathogen response, the long duration of IKK/NF- κ B activity induced by LPS allows for induction genes that are poorly induced by transient TNF stimulation (Werner et al., 2005). A recent study quantified the contribution of long mRNA half-life to decoding differential duration of NF- κ B activity for several dozen genes, but also showed that chromatin-associated mechanisms play important roles as well (Sen et al., 2020).

Nucleosome displacement also constitutes a kinetically “slow” step such that signal-dependent transcription factors (SDTF) binding to promoters or enhancers (Kaikkonen et al., 2013; Tong et al., 2016), co-activator recruitment, and pre-initiation complex (PIC) assembly and polymerase recruitment (Bhaumik, 2011) may, in principle, decode temporal dynamics of signaling. For example, LPS-induced NF- κ B target genes fall into two categories: those with open chromatin immediately available for NF- κ B binding and ready transcriptional induction, and those




Figure 4. Mechanisms that decode combinatorial and temporal codes of pathway activity to achieve stimulus-specific responses.

(A) Decoding of pathway-specific effector combinations occurs through sequential logic gates that involve pre-transcriptional mechanisms, such as (left top) NF-κB and ISGF3 forming a sequential OR gate, allowing sustained responses; (left bottom) IRF3-mediated nucleosome movement for *Ccl5*, allowing NF-κB binding and forming a sequential AND gate; and (right) post-transcriptional mechanisms combining NF-κB-induced transcription AND MAPK38 processing and stabilization of *Tnf* mRNA.

(B) Coherent feedforward loops, where one regulator positively controls a target and another positive regulator. Left: NF-κB induces transcription of *Cebpd*, which activates a second wave of response genes controlled by both NF-κB and CEPBδ. Here, coherent FFLs combined with AND gates allow duration decoding. Right: similarly, IRF3-induced *lfnb1* generates a coherent feedforward loop through activation of IFNAR and subsequently ISGF3, which binds similar gene promoters as IRF3.

(C) Incoherent feedforward loops, where one regulator positively controls a target as well as a negative regulator. Together with combinatorial decoding principles they allow for fold-change detection and mediate biphasic responses. Left: NF-κB-induced expression of the competing p50:p50 homodimer provides memory of previous NF-κB abundances for fold-change detection on NF-κB target genes such as *I/8*. Right: NF-κB-induced transcription of *Atf3*, which in turn represses NF-κB target genes. This results in the diminishing of the second phase of response gene expression when NF-κB activity is high. Corner arrow indicates active transcription.

requiring chromatin remodeling before NF-κB can bind (Kayama et al., 2008; Natoli, 2009; Ramirez-Carrozzi et al., 2009; Saccani et al., 2001). One recent study quantified the contribution of chromatin-associated mechanisms for differentiating between LPS-versus TNF-induced (2–3 h versus 1 h) NF-κB activity for 81 immune response genes (Sen et al., 2020). Interestingly, the chromatin-associated mechanisms provided a higher degree of specificity than the aforementioned long mRNA half-life mechanism and pertained particularly to immune effectors such as *Mmp3*, *Pfr1*, *Pilra*, and also *Ccl5*.

The formation of *de novo* enhancers is a third kinetically slow step that can decode temporal dynamics, but it is less reversible than the two previous examples. While poised target gene expression did not markedly differ in cells with oscillatory versus non-oscillatory NF-κB activity (Barken et al., 2005), a recent study

multi-step unwrapping and rewinding dynamics of nucleosomal DNA-histone interactions, displacing the nucleosome and allowing the deposition of histone marks. A negative feedback mutant that converted normally oscillatory TNF-induced NF-κB activity to non-oscillatory activity allowed for more efficient nucleosome eviction and increased chromatin accessibility, and the subsequent formation of *de novo* enhancers enabled novel gene expression programs upon a secondary stimulation. SDTF dynamics may thus also be a critical determinant for epigenomic reprogramming that may allow immune sentinel cells to function in an exposure-history-dependent manner, potentially conferring innate immune memory (Netea et al., 2016).

A gene regulatory network where an effector positively controls a target both directly and indirectly via another positive regulator forms a coherent feedforward loop. Coherent

feedforward loops, which comprise of an AND logic gate, may decode differential TF durations (Figure 4B). For example, IL-6 expression was found to depend not only on NF- κ B, but also the secondary TF C/EBP δ (Litvak et al., 2009). Because NF- κ B activity induces the expression of C/EBP δ , they form a coherent feedforward loop to activate IL-6. However, a short exposure to LPS will not provide long-lasting NF- κ B that can synergize with *de novo* C/EBP δ , thus preventing induction of secondary IL-6-mediated inflammation. The regulatory relationship of IRF3 and ISGF3 may also constitute a coherent feedforward loop on genes that require both IRF3 and ISGF3 for full activation. If, for example, IRF3 triggers a key chromatin remodeling step that then allows ISGF3 to bind and activate transcription, such target genes would primarily be induced in the infected (pathogen-exposed) cell rather than the bystander cells. A transcriptomic analysis yielded just a few candidates (e.g., *Cxcl10*, *Iffit3*) for such a regulatory mechanism (Ourthiague et al., 2015) and whether they differentiate IRF3-ISGF3 temporal features was not explored.

Another type of regulatory network motif, in which a signaling effector positively controls a target as well as a negative regulator of the same target, is referred to as an incoherent feedforward loop. Incoherent feedforward loops provide decoding mechanism for differences in fold changes (Figure 4C). For example, the decoding of fold changes of NF- κ B activity on a set of target genes was reported to be governed by an incoherent feedforward loop involving the activation-domain lacking p50:p50 homodimer (Lee et al., 2014). A subset of TNF-induced genes was found to correlate particularly well with NF- κ B fold changes in single cells. Interestingly, p50:p50 homodimers were also shown to bind a subset of ISREs and thus tune the responsiveness of ISGF3 target genes (Cheng et al., 2011). In an earlier example, an incoherent feedforward loop was shown to prevent run-away expression of inflammatory cytokines (Gilchrist et al., 2006). ATF3 was found to repress the NF- κ B-induced expression of IL-6 and IL-12B. As ATF3 itself is induced by NF- κ B, its repressive role on IL-6 and IL-12B becomes relevant when NF- κ B activity is high and sustained, thereby preventing over-production of these potent inflammatory cytokines.

In further illustration of the precise temporal regulation of gene expression, incoherent feed forward loops may also involve post-transcriptional mechanisms. MicroRNAs known as inflam-mamIRs, mir155 and mir146, are NF- κ B inducible and target immune response mRNAs (Rothchild et al., 2016; Testa et al., 2017). Mir146 targets regulators of the TLR signaling pathway and this repressive function serves to fine tune the innate immune response, whereas mir155 acts in an opposing and dominant function to further potentiate NF- κ B response (Mann et al., 2017). Similarly, the RNA editing protein ADAR1 provides incoherent feedforward control because it is induced by IFN and then acts to repress RLR signaling, dampening the IFN response and preventing autoimmunity (Lamers et al., 2019; Mannion et al., 2014). While these regulatory circuits have the potential to decode dynamic features of the pathogen-responsive temporal code, their true functional impact will need to be addressed in future studies.

Decoding mechanisms are intricately tied to immune sentinel functions of context dependence and memory. Beyond generating the stimulus-specificity of immediate gene expression

responses, decoding mechanisms that invoke chromatin remodeling are particularly important to the establishment of stimulus-specific innate immune memory. The eviction of nucleosomes and the deposition of new histone marks are a downstream consequence of stimulus-specific chromatin remodeling, though the functional and physiological consequences of this reprogramming, especially in disease contexts, remains to be explored.

CONCLUDING REMARKS

Recent work has begun to document the capacity of immune sentinel cells to mount pathogen-specific responses, as well as provide a set of mechanistic studies to support the notions of combinatorial and temporal coding. These insights may serve as a basis for addressing key questions through future studies.

First, while the mechanisms underlying temporal coding in the NF- κ B signaling module are relatively well understood, the integration of combinatorial and temporal signaling via coordinated activities of IRF, JNK, and MAPKp38 pathways is a next big challenge in the immune signaling field. Temporal dynamics and pathway branching of signaling initiated by cytosolic PRRs such as cGAS, RLRs, and NLRs are also understudied.

Second, understanding how combinatorial and temporal signaling codes are interpreted to produce gene expression requires substantial effort, in part because every gene is controlled by a seemingly unique gene regulatory network. To study the connection between encoding and decoding of signaling activity, it will be necessary to develop efficient workflows that directly connect temporal effector activity profiles, ideally for multiple effectors, with gene expression or cytokine secretion measurements in the same cells (Gutschow et al., 2019; Lane et al., 2017).

Third, and related, at the single-cell level, both stimulus-responsive signaling activity and gene expression are highly heterogeneous, which may be essential for the breadth, robustness, and adaptability of the immune response but may also contribute to pathology (Satija and Shalek, 2014). While live cell imaging has been successful in characterizing the cell-to-cell variability of signaling activity, the stochasticity of decoding mechanisms at the gene promoter likely plays an especially important role in determining gene expression responses of a specific cell. This requires innovative integration of live cell imaging and necessarily destructive single-cell technologies, such as single-cell RNA sequencing (scRNA-seq), single-molecule fluorescence in-situ hybridization (smFISH), single-cell assay for transposase-accessible chromatin using sequencing (scATAC-seq), or single-cell chromatin immuno-precipitation followed by sequencing (scChIP-seq) to probe expression of native chromatin and endogenous genes (rather than ectopic reporters), via data driven and mechanistic modeling approaches.

Fourth, with such workflows in place, future work will be able to explore how encoding and decoding mechanisms are a function of microenvironmental context and the cell's exposure history. Cytokines in the microenvironment, which guide the functional specialization of immune sentinel cells, may alter the encoding of combinatorial and temporal signaling profiles, while exposure history that changes the cell's baseline epigenome affects how signals are decoded (Cheng et al., 2019; Kang et al., 2017; Park et al., 2017).

Detailed and predictive knowledge of how immune sentinel responses are regulated will undoubtedly impact our understanding of a range of immune-related disease, whether immunopathological responses to infectious disease (e.g., as seen during severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2] infections) or autoinflammatory and autoimmune disorders (Funes et al., 2018; Gustine and Jones, 2021). As pathogenesis may be triggered by small subsets of outlier cells, understanding the basis for the heterogeneity of responses at the single-cell level is key. By characterizing macrophages derived from a Sjögren's syndrome mouse model, a recent study revealed that a minority of cells show stimulus-confused NF- κ B dynamics and gene expression (Adelaja et al., 2021). We therefore suggest that a quantitative and mechanistic understanding of immune sentinel responses will enable a better characterization and diagnosis of auto-inflammatory and auto-immune diseases and support the development of therapeutic strategies aimed at correcting disease-associated miscoding.

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.immuni.2021.08.018>.

ACKNOWLEDGMENTS

S.L. is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 419234150. K.M.S. is supported by the UCLA Medical Scientist Training Program (NIH NIGMS T32 GM008042). Research in the Hoffmann lab on described topics is funded by NIH R01AI127867, R01AI132835, R01AI127864, and R01GM117134.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Abe, T., and Barber, G.N. (2014). Cytosolic-DNA-mediated, STING-dependent proinflammatory gene induction necessitates canonical NF- κ B activation through TBK1. *J. Virol.* *88*, 5328–5341.

Abe, H., Satoh, J., Shirasaka, Y., Kogure, A., Kato, H., Ito, S., and Fujita, T. (2020). Priming Phosphorylation of TANK-Binding Kinase 1 by I κ B Kinase β Is Essential in Toll-Like Receptor 3/4 Signaling. *Mol. Cell. Biol.* *40*, 40.

Ablasser, A., and Chen, Z.J. (2019). cGAS in action: Expanding roles in immunity and inflammation. *Science* *363*, eaat8657.

Ablasser, A., and Hur, S. (2020). Regulation of cGAS- and RLR-mediated immunity to nucleic acids. *Nat. Immunol.* *21*, 17–29.

Adelaja, A., Taylor, B., Sheu, K.M., Liu, Y., Luecke, S., and Hoffmann, A. (2021). Six distinct NF κ B signaling codons convey discrete information to distinguish stimuli and enable appropriate macrophage responses. *Immunity* *54*, 916–930.

Aikin, T.J., Peterson, A.F., Pokrass, M.J., Clark, H.R., and Regot, S. (2019). Collective MAPK Signaling Dynamics Coordinates Epithelial Homeostasis. *bioRxiv*, 826917.

Albeck, J.G., Burke, J.M., Spencer, S.L., Lauffenburger, D.A., and Sorger, P.K. (2008a). Modeling a snap-action, variable-delay switch controlling extrinsic cell death. *PLoS Biol.* *6*, 2831–2852.

Albeck, J.G., Burke, J.M., Aldridge, B.B., Zhang, M., Lauffenburger, D.A., and Sorger, P.K. (2008b). Quantitative analysis of pathways controlling extrinsic apoptosis in single cells. *Mol. Cell* *30*, 11–25.

Altan-Bonnet, G., and Germain, R.N. (2005). Modeling T cell antigen discrimination based on feedback control of digital ERK responses. *PLoS Biol.* *3*, e356.

Amir, I., Garber, M., Chevrier, N., Leite, A.P., Donner, Y., Eisenhaure, T., Guttman, M., Grenier, J.K., Li, W., Zuk, O., et al. (2009). Unbiased reconstruction of a mammalian transcriptional network mediating pathogen responses. *Science* *326*, 257–263.

Ashall, L., Horton, C.A., Nelson, D.E., Paszek, P., Harper, C.V., Sillitoe, K., Ryan, S., Spiller, D.G., Unitt, J.F., Broomhead, D.S., et al. (2009). Pulsatile stimulation determines timing and specificity of NF- κ B-dependent transcription. *Science* *324*, 242–246.

Bagnall, J., Boddington, C., England, H., Brignall, R., Downton, P., Alsoufi, Z., Boyd, J., Rowe, W., Bennett, A., Walker, C., et al. (2018). Quantitative analysis of competitive cytokine signaling predicts tissue thresholds for the propagation of macrophage activation. *Sci. Signal.* *11*, eaaf3998.

Balka, K.R., and De Nardo, D. (2019). Understanding early TLR signaling through the Myddosome. *J. Leukoc. Biol.* *105*, 339–351.

Bardwell, L. (2008). Signal transduction: turning a switch into a rheostat. *Curr. Biol.* *18*, R910–R912.

Barken, D., Wang, C.J., Kearns, J., Cheong, R., Hoffmann, A., and Levchenko, A. (2005). Comment on "Oscillations in NF- κ B signaling control the dynamics of gene expression". *Science* *308*, 52, 52, author reply 52.

Barrat, F.J., Crow, M.K., and Ivashkiv, L.B. (2019). Interferon target-gene expression and epigenomic signatures in health and disease. *Nat. Immunol.* *20*, 1574–1583.

Becker, V., Schilling, M., Bachmann, J., Baumann, U., Raue, A., Maiwald, T., Timmer, J., and Klingmüller, U. (2010). Covering a broad dynamic range: information processing at the erythropoietin receptor. *Science* *328*, 1404–1408.

Behar, M., and Hoffmann, A. (2010). Understanding the temporal codes of intra-cellular signals. *Curr. Opin. Genet. Dev.* *20*, 684–693.

Behar, M., and Hoffmann, A. (2013). Tunable signal processing through a kinase control cycle: the IKK signaling node. *Biophys. J.* *105*, 231–241.

Beinke, S., Robinson, M.J., Hugunin, M., and Ley, S.C. (2004). Lipopolysaccharide activation of the TPL-2/MEK/extracellular signal-regulated kinase mitogen-activated protein kinase cascade is regulated by I κ B kinase-induced proteolysis of NF- κ B1 p105. *Mol. Cell. Biol.* *24*, 9658–9667.

Bhaumik, S.R. (2011). Distinct regulatory mechanisms of eukaryotic transcriptional activation by SAGA and TFIID. *Biochim. Biophys. Acta* *1809*, 97–108.

Boraschi, D., Italiani, P., Weil, S., and Martin, M.U. (2018). The family of the interleukin-1 receptors. *Immunol. Rev.* *281*, 197–232.

Bray, D., Levin, M.D., and Morton-Firth, C.J. (1998). Receptor clustering as a cellular mechanism to control sensitivity. *Nature* *393*, 85–88.

Brignall, R., Moody, A.T., Mathew, S., and Gaudet, S. (2019). Considering Abundance, Affinity, and Binding Site Availability in the NF- κ B Target Selection Puzzle. *Front. Immunol.* *10*, 609.

Brondello, J.-M., Pouyssegur, J., and McKenzie, F.R. (1999). Reduced MAP kinase phosphatase-1 degradation after p42/p44MAPK-dependent phosphorylation. *Science* *286*, 2514–2517.

Buchler, N.E., Gerland, U., and Hwa, T. (2003). On schemes of combinatorial transcription logic. *Proc. Natl. Acad. Sci. USA* *100*, 5136–5141.

Cadena, C., Ahmad, S., Xavier, A., Willemsen, J., Park, S., Park, J.W., Oh, S.-W., Fujita, T., Hou, F., Binder, M., and Hur, S. (2019). Ubiquitin-Dependent and -Independent Roles of E3 Ligase RIPLET in Innate Immunity. *Cell* *177*, 1187–1200.

Caldwell, A.B., Cheng, Z., Vargas, J.D., Birnbaum, H.A., and Hoffmann, A. (2014). Network dynamics determine the autocrine and paracrine signaling functions of TNF. *Genes Dev.* *28*, 2120–2133.

Caré, B.R., and Soula, H.A. (2011). Impact of receptor clustering on ligand binding. *BMC Syst. Biol.* *5*, 48.

Cargnello, M., and Roux, P.P. (2011). Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiol. Mol. Biol. Rev.* *75*, 50–83.

Caruso, R., Warner, N., Inohara, N., and Núñez, G. (2014). NOD1 and NOD2: signaling, host defense, and inflammatory disease. *Immunity* *41*, 898–908.

- Catrysse, L., Vereecke, L., Beyaert, R., and van Loo, G. (2014). A20 in inflammation and autoimmunity. *Trends Immunol.* **35**, 22–31.
- Chakraborty, A.K., and Weiss, A. (2014). Insights into the initiation of TCR signaling. *Nat. Immunol.* **15**, 798–807.
- Chen, H., and Jiang, Z. (2013). The essential adaptors of innate immune signaling. *Protein Cell* **4**, 27–39.
- Chen, J., Almo, S.C., and Wu, Y. (2017). General principles of binding between cell surface receptors and multi-specific ligands: A computational study. *PLoS Comput. Biol.* **13**, e1005805.
- Cheng, C.S., Feldman, K.E., Lee, J., Verma, S., Huang, D.B., Huynh, K., Chang, M., Ponomarenko, J.V., Sun, S.C., Benedict, C.A., et al. (2011). The specificity of innate immune responses is enforced by repression of interferon response elements by NF- κ B p50. *Sci. Signal.* **4**, ra11.
- Cheng, Z., Taylor, B., Ourthiague, D.R., and Hoffmann, A. (2015). Distinct single-cell signaling characteristics are conferred by the MyD88 and TRIF pathways during TLR4 activation. *Sci. Signal.* **8**, ra69.
- Cheng, C.S., Behar, M.S., Suryawanshi, G.W., Feldman, K.E., Spreafico, R., and Hoffmann, A. (2017). Iterative Modeling Reveals Evidence of Sequential Transcriptional Control Mechanisms. *Cell Syst.* **4**, 330–343.
- Cheng, Q., Behzadi, F., Sen, S., Ohta, S., Spreafico, R., Teles, R., Modlin, R.L., and Hoffmann, A. (2019). Sequential conditioning-stimulation reveals distinct gene- and stimulus-specific effects of Type I and II IFN on human macrophage functions. *Sci. Rep.* **9**, 5288.
- Cheng, Q.J., Ohta, S., Sheu, K.M., Spreafico, R., Adelaja, A., Taylor, B., and Hoffmann, A. (2021). NF- κ B dynamics determine the stimulus specificity of epigenomic reprogramming in macrophages. *Science* **372**, 1349–1353.
- Cheong, R., Bergmann, A., Werner, S.L., Regal, J., Hoffmann, A., and Levchenko, A. (2006). Transient I κ B kinase activity mediates temporal NF- κ B dynamics in response to a wide range of tumor necrosis factor- α doses. *J. Biol. Chem.* **281**, 2945–2950.
- Cheong, R., Rhee, A., Wang, C.J., Nemenman, I., and Levchenko, A. (2011). Information transduction capacity of noisy biochemical signaling networks. *Science* **334**, 354–358.
- Chhabra, S., Liu, L., Goh, R., Kong, X., and Warmflash, A. (2019). Dissecting the dynamics of signaling events in the BMP, WNT, and NODAL cascade during self-organized fate patterning in human gastruloids. *PLoS Biol.* **17**, e3000498.
- Chopra, S.S., Jenney, A., Palmer, A., Niepel, M., Chung, M., Mills, C., Sivakumaran, S.C., Liu, Q., Chen, J.-Y., Yapp, C., et al. (2020). Torin2 Exploits Replication and Checkpoint Vulnerabilities to Cause Death of PI3K-Activated Triple-Negative Breast Cancer Cells. *Cell Syst.* **10**, 66–81.
- Cohen, P., and Strickson, S. (2017). The role of hybrid ubiquitin chains in the MyD88 and other innate immune signalling pathways. *Cell Death Differ.* **24**, 1153–1159.
- Covert, M.W., Leung, T.H., Gaston, J.E., and Baltimore, D. (2005). Achieving stability of lipopolysaccharide-induced NF- κ B activation. *Science* **309**, 1854–1857.
- Cusson-Hermance, N., Khurana, S., Lee, T.H., Fitzgerald, K.A., and Kelliher, M.A. (2005). Rip1 mediates the Trif-dependent toll-like receptor 3- and 4-induced NF- κ B activation but does not contribute to interferon regulatory factor 3 activation. *J. Biol. Chem.* **280**, 36560–36566.
- De, A., Dainichi, T., Rathinam, C.V., and Ghosh, S. (2014). The deubiquitinase activity of A20 is dispensable for NF- κ B signaling. *EMBO Rep.* **15**, 775–783.
- de Oliveira Mann, C.C., Orzalli, M.H., King, D.S., Kagan, J.C., Lee, A.S.Y., and Kranzusch, P.J. (2019). Modular Architecture of the STING C-Terminal Tail Allows Interferon and NF- κ B Signaling Adaptation. *Cell Rep.* **27**, 1165–1175.
- de Weerd, N.A., and Nguyen, T. (2012). The interferons and their receptors—distribution and regulation. *Immunol. Cell Biol.* **90**, 483–491.
- Deguine, J., and Barton, G.M. (2014). MyD88: a central player in innate immune signaling. *F1000Prime Rep.* **6**, 97.
- Delhase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999). Positive and negative regulation of I κ B kinase activity through IKK β subunit phosphorylation. *Science* **284**, 309–313.
- Dostert, C., Grusdat, M., Letellier, E., and Brenner, D. (2019). The TNF Family of Ligands and Receptors: Communication Modules in the Immune System and Beyond. *Physiol. Rev.* **99**, 115–160.
- Doyle, S., Vaidya, S., O’Connell, R., Dadgostar, H., Dempsey, P., Wu, T., Rao, G., Sun, R., Haberland, M., Modlin, R., and Cheng, G. (2002). IRF3 mediates a TLR3/TLR4-specific antiviral gene program. *Immunity* **17**, 251–263.
- Dunphy, G., Flannery, S.M., Almine, J.F., Connolly, D.J., Paulus, C., Jönsson, K.L., Jakobsen, M.R., Nevels, M.M., Bowie, A.G., and Unterholzner, L. (2018). Non-canonical Activation of the DNA Sensing Adaptor STING by ATM and IFI16 Mediates NF- κ B Signaling after Nuclear DNA Damage. *Mol. Cell* **71**, 745–760.
- Ea, C.-K., Deng, L., Xia, Z.-P., Pineda, G., and Chen, Z.J. (2006). Activation of IKK by TNF α requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol. Cell* **22**, 245–257.
- Emmerich, C.H., Ordureau, A., Strickson, S., Arthur, J.S.C., Pedrioli, P.G.A., Komander, D., and Cohen, P. (2013). Activation of the canonical IKK complex by K63/M1-linked hybrid ubiquitin chains. *Proc. Natl. Acad. Sci. USA* **110**, 15247–15252.
- Eng, C.L., Lawson, M., Zhu, Q., Dries, R., Koulens, N., Takei, Y., Yun, J., Cronin, C., Karp, C., Yuan, G.-C., and Cai, L. (2019). Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. *Nature* **568**, 235–239.
- Fang, R., Jiang, Q., Zhou, X., Wang, C., Guan, Y., Tao, J., Xi, J., Feng, J.-M., and Jiang, Z. (2017a). MAVS activates TBK1 and IKK ϵ through TRAFs in NEMO dependent and independent manner. *PLoS Pathog.* **13**, e1006720.
- Fang, R., Wang, C., Jiang, Q., Lv, M., Gao, P., Yu, X., Mu, P., Zhang, R., Bi, S., Feng, J.M., et al. (2017b). NEMO-IKK β Are Essential for IRF3 and NF- κ B Activation in the cGAS-STING Pathway. *J. Immunol.* **199**, 3222–3233.
- Feinerman, O., Germain, R.N., and Altan-Bonnet, G. (2008a). Quantitative challenges in understanding ligand discrimination by α beta T cells. *Mol. Immunol.* **45**, 619–631.
- Feinerman, O., Veiga, J., Dorfman, J.R., Germain, R.N., and Altan-Bonnet, G. (2008b). Variability and robustness in T cell activation from regulated heterogeneity in protein levels. *Science* **321**, 1081–1084.
- Ferraro, R., Zhou, H., Shan, Y., Liu, Q., Li, Q., Shaw, D.E., Li, X., and Wu, H. (2014). IRAK4 dimerization and trans-autophosphorylation are induced by Myddosome assembly. *Mol. Cell* **55**, 891–903.
- Ferrell, J.E., Jr., and Bhatt, R.R. (1997). Mechanistic studies of the dual phosphorylation of mitogen-activated protein kinase. *J. Biol. Chem.* **272**, 19008–19016.
- Ferrell, J.E., Jr., and Ha, S.H. (2014). Ultrasensitivity part III: cascades, bistable switches, and oscillators. *Trends Biochem. Sci.* **39**, 612–618.
- Foreman, R., and Wollman, R. (2020). Mammalian gene expression variability is explained by underlying cell state. *Mol. Syst. Biol.* **16**, e9146.
- Freaney, J.E., Kim, R., Mandhana, R., and Horvath, C.M. (2013). Extensive cooperation of immune master regulators IRF3 and NF- κ B in RNA Pol II recruitment and pause release in human innate antiviral transcription. *Cell Rep.* **4**, 959–973.
- Fric, J., Zelante, T., Wong, A.Y.W., Mertes, A., Yu, H.-B., and Ricciardi-Castagnoli, P. (2012). NFAT control of innate immunity. *Blood* **120**, 1380–1389.
- Friedman, C.S., O’Donnell, M.A., Legarda-Addison, D., Ng, A., Cárdenas, W.B., Yount, J.S., Moran, T.M., Basler, C.F., Komuro, A., Horvath, C.M., et al. (2008). The tumour suppressor CYLD is a negative regulator of RIG-I-mediated antiviral response. *EMBO Rep.* **9**, 930–936.
- Funami, K., Sasai, M., Oshiumi, H., Seya, T., and Matsumoto, M. (2008). Homo-oligomerization is essential for Toll/interleukin-1 receptor domain-containing adaptor molecule-1-mediated NF- κ B and interferon regulatory factor-3 activation. *J. Biol. Chem.* **283**, 18283–18291.
- Funes, S.C., Rios, M., Escobar-Vera, J., and Kalergis, A.M. (2018). Implications of macrophage polarization in autoimmunity. *Immunology* **154**, 186–195.
- Ganti, R.S., Lo, W.-L., McAfee, D.B., Groves, J.T., Weiss, A., and Chakraborty, A.K. (2020). How the T cell signaling network processes information to discriminate between self and agonist ligands. *Proc. Natl. Acad. Sci. USA* **117**, 26020–26030.

- Gantke, T., Sriskantharajah, S., Sadowski, M., and Ley, S.C. (2012). I κ B kinase regulation of the TPL-2/ERK MAPK pathway. *Immunol. Rev.* *246*, 168–182.
- Gaudet, S., Spencer, S.L., Chen, W.W., and Sorger, P.K. (2012). Exploring the contextual sensitivity of factors that determine cell-to-cell variability in receptor-mediated apoptosis. *PLoS Comput. Biol.* *8*, e1002482.
- Gilchrist, M., Thorsson, V., Li, B., Rust, A.G., Korb, M., Roach, J.C., Kennedy, K., Hai, T., Bolouri, H., and Aderem, A. (2006). Systems biology approaches identify ATF3 as a negative regulator of Toll-like receptor 4. *Nature* *441*, 173–178.
- Gottschalk, R.A., Martins, A.J., Angermann, B.R., Dutta, B., Ng, C.E., Uderhardt, S., Tsang, J.S., Fraser, I.D., Meier-Schellersheim, M., and Germain, R.N. (2016). Distinct NF- κ B and MAPK Activation Thresholds Uncouple Steady-State Microbe Sensing from Anti-pathogen Inflammatory Responses. *Cell Syst.* *2*, 378–390.
- Gottschalk, R.A., Dorrington, M.G., Dutta, B., Krauss, K.S., Martins, A.J., Uderhardt, S., Chan, W., Tsang, J.S., Torabi-Parizi, P., Fraser, I.D., and Germain, R.N. (2019). IFN-mediated negative feedback supports bacteria class-specific macrophage inflammatory responses. *eLife* *8*, e46836.
- Gu, M., Zhang, T., Lin, W., Liu, Z., Lai, R., Xia, D., Huang, H., and Wang, X. (2014). Protein phosphatase PP1 negatively regulates the Toll-like receptor and RIG-I-like receptor-triggered production of type I interferon by inhibiting IRF3 phosphorylation at serines 396 and 385 in macrophage. *Cell. Signal.* *26*, 2930–2939.
- Gustine, J.N., and Jones, D. (2021). Immunopathology of Hyperinflammation in COVID-19. *Am. J. Pathol.* *191*, 4–17.
- Gutschow, M.V., Mason, J.C., Lane, K.M., Maayan, I., Hughey, J.J., Bajar, B.T., Amatya, D.N., Valle, S.D., and Covert, M.W. (2019). Combinatorial processing of bacterial and host-derived innate immune stimuli at the single-cell level. *Mol. Biol. Cell* *30*, 282–292.
- Haas, T.L., Emmerich, C.H., Gerlach, B., Schmukle, A.C., Cordier, S.M., Rieser, E., Feltham, R., Vince, J., Warnken, U., Wenger, T., et al. (2009). Recruitment of the linear ubiquitin chain assembly complex stabilizes the TNF-R1 signaling complex and is required for TNF-mediated gene induction. *Mol. Cell* *36*, 831–844.
- Häcker, H., and Karin, M. (2006). Regulation and function of IKK and IKK-related kinases. *Sci. STKE* *2006*, re13.
- Hao, S., and Baltimore, D. (2009). The stability of mRNA influences the temporal order of the induction of genes encoding inflammatory molecules. *Nat. Immunol.* *10*, 281–288.
- Hayden, M.S., and Ghosh, S. (2004). Signaling to NF- κ B. *Genes Dev.* *18*, 2195–2224.
- Heemskerck, I., Burt, K., Miller, M., Chhabra, S., Guerra, M.C., Liu, L., and Warmflash, A. (2019). Rapid changes in morphogen concentration control self-organized patterning in human embryonic stem cells. *eLife* *8*, e40526.
- Heinrich, R., Neel, B.G., and Rapoport, T.A. (2002). Mathematical models of protein kinase signal transduction. *Mol. Cell* *9*, 957–970.
- Hoffmann, A. (2016). Immune Response Signaling: Combinatorial and Dynamic Control. *Trends Immunol.* *37*, 570–572.
- Hoffmann, A., and Baltimore, D. (2006). Circuitry of nuclear factor κ B signaling. *Immunol. Rev.* *210*, 171–186.
- Hoffmann, A., Levchenko, A., Scott, M.L., and Baltimore, D. (2002). The I κ B-NF- κ B signaling module: temporal control and selective gene activation. *Science* *298*, 1241–1245.
- Honda, K., Yanai, H., Mizutani, T., Negishi, H., Shimada, N., Suzuki, N., Ohba, Y., Takaoka, A., Yeh, W.-C., and Taniguchi, T. (2004). Role of a transcriptional-transcriptional processor complex involving MyD88 and IRF-7 in Toll-like receptor signaling. *Proc. Natl. Acad. Sci. USA* *101*, 15416–15421.
- Honda, K., Ohba, Y., Yanai, H., Negishi, H., Mizutani, T., Takaoka, A., Taya, C., and Taniguchi, T. (2005). Spatiotemporal regulation of MyD88-IRF-7 signalling for robust type-I interferon induction. *Nature* *434*, 1035–1040.
- Honda, K., Takaoka, A., and Taniguchi, T. (2006). Type I interferon [corrected] gene induction by the interferon regulatory factor family of transcription factors. *Immunity* *25*, 349–360.
- Hoshino, K., Sugiyama, T., Matsumoto, M., Tanaka, T., Saito, M., Hemmi, H., Ohara, O., Akira, S., and Kaisho, T. (2006). I κ B kinase- α is critical for interferon- α production induced by Toll-like receptors 7 and 9. *Nature* *440*, 949–953.
- Hu, H., and Sun, S.-C. (2016). Ubiquitin signaling in immune responses. *Cell Res.* *26*, 457–483.
- Huang, C.Y., and Ferrell, J.E., Jr. (1996). Ultrasensitivity in the mitogen-activated protein kinase cascade. *Proc. Natl. Acad. Sci. USA* *93*, 10078–10083.
- Hughey, J.J., Gutschow, M.V., Bajar, B.T., and Covert, M.W. (2015). Single-cell variation leads to population invariance in NF- κ B signaling dynamics. *Mol. Biol. Cell* *26*, 583–590.
- Husebye, H., Halaas, Ø., Stenmark, H., Tunheim, G., Sandanger, Ø., Bogen, B., Brech, A., Latz, E., and Espevik, T. (2006). Endocytic pathways regulate Toll-like receptor 4 signaling and link innate and adaptive immunity. *EMBO J.* *25*, 683–692.
- Ishitani, T., Takaesu, G., Ninomiya-Tsuji, J., Shibuya, H., Gaynor, R.B., and Matsumoto, K. (2003). Role of the TAB2-related protein TAB3 in IL-1 and TNF signaling. *EMBO J.* *22*, 6277–6288.
- Ivashkiv, L.B., and Donlin, L.T. (2014). Regulation of type I interferon responses. *Nat. Rev. Immunol.* *14*, 36–49.
- Janeway, C.A., Jr., and Medzhitov, R. (2002). Innate immune recognition. *Annu. Rev. Immunol.* *20*, 197–216.
- Jiang, Z., Mak, T.W., Sen, G., and Li, X. (2004). Toll-like receptor 3-mediated activation of NF- κ B and IRF3 diverges at Toll-IL-1 receptor domain-containing adapter inducing IFN- β . *Proc. Natl. Acad. Sci. USA* *101*, 3533–3538.
- Junkin, M., Kaestli, A.J., Cheng, Z., Jordi, C., Albayrak, C., Hoffmann, A., and Tay, S. (2016). High-Content Quantification of Single-Cell Immune Dynamics. *Cell Rep.* *15*, 411–422.
- Kagan, J.C., Su, T., Horng, T., Chow, A., Akira, S., and Medzhitov, R. (2008). TRAM couples endocytosis of Toll-like receptor 4 to the induction of interferon- β . *Nat. Immunol.* *9*, 361–368.
- Kaikkonen, M.U., Spann, N.J., Heinz, S., Romanoski, C.E., Allison, K.A., Stender, J.D., Chun, H.B., Tough, D.F., Prinjha, R.K., Benner, C., and Glass, C.K. (2013). Remodeling of the enhancer landscape during macrophage activation is coupled to enhancer transcription. *Mol. Cell* *51*, 310–325.
- Kanayama, A., Seth, R.B., Sun, L., Ea, C.-K., Hong, M., Shaito, A., Chiu, Y.-H., Deng, L., and Chen, Z.J. (2004). TAB2 and TAB3 activate the NF- κ B pathway through binding to polyubiquitin chains. *Mol. Cell* *15*, 535–548.
- Kang, K., Park, S.H., Chen, J., Qiao, Y., Giannopoulou, E., Berg, K., Hanidu, A., Li, J., Nabozny, G., Kang, K., et al. (2017). Interferon- γ Represses M2 Gene Expression in Human Macrophages by Disassembling Enhancers Bound by the Transcription Factor MAF. *Immunity* *47*, 235–250.
- Kawai, T., and Akira, S. (2011). Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity* *34*, 637–650.
- Kawai, T., Sato, S., Ishii, K.J., Coban, C., Hemmi, H., Yamamoto, M., Terai, K., Matsuda, M., Inoue, J., Uematsu, S., et al. (2004). Interferon- α induction through Toll-like receptors involves a direct interaction of IRF7 with MyD88 and TRAF6. *Nat. Immunol.* *5*, 1061–1068.
- Kayama, H., Ramirez-Carrozzi, V.R., Yamamoto, M., Mizutani, T., Kuwata, H., Iba, H., Matsumoto, M., Honda, K., Smale, S.T., and Takeda, K. (2008). Class-specific regulation of pro-inflammatory genes by MyD88 pathways and I κ B-zeta. *J. Biol. Chem.* *283*, 12468–12477.
- Kearns, J.D., Basak, S., Werner, S.L., Huang, C.S., and Hoffmann, A. (2006). I κ B β provides negative feedback to control NF- κ B oscillations, signaling dynamics, and inflammatory gene expression. *J. Cell Biol.* *173*, 659–664.
- Kellogg, R.A., and Tay, S. (2015). Noise facilitates transcriptional control under dynamic inputs. *Cell* *160*, 381–392.
- Kensche, T., Tokunaga, F., Ikeda, F., Goto, E., Iwai, K., and Dikic, I. (2012). Analysis of nuclear factor- κ B (NF- κ B) essential modulator (NEMO) binding to linear and lysine-linked ubiquitin chains and its role in the activation of NF- κ B. *J. Biol. Chem.* *287*, 23626–23634.

- Kholodenko, B.N., and Birtwistle, M.R. (2009). Four-dimensional dynamics of MAPK information processing systems. *Wiley Interdiscip. Rev. Syst. Biol. Med.* *1*, 28–44.
- Krebs, D.L., and Hilton, D.J. (2001). SOCS proteins: negative regulators of cytokine signaling. *Stem Cells* *19*, 378–387.
- Kundert, K., Lucas, J.E., Watters, K.E., Fellmann, C., Ng, A.H., Heineke, B.M., Fitzsimmons, C.M., Oakes, B.L., Qu, J., Prasad, N., et al. (2019). Controlling CRISPR-Cas9 with ligand-activated and ligand-deactivated sgRNAs. *Nat. Commun.* *10*, 2127.
- Kuwano, Y., Kim, H.H., Abdelmohsen, K., Pullmann, R., Jr., Martindale, J.L., Yang, X., and Gorospe, M. (2008). MKP-1 mRNA stabilization and translational control by RNA-binding proteins HuR and NF90. *Mol. Cell. Biol.* *28*, 4562–4575.
- Lamers, M.M., van den Hoogen, B.G., and Haagmans, B.L. (2019). ADAR1: “Editor-in-Chief” of Cytoplasmic Innate Immunity. *Front. Immunol.* *10*, 1763.
- Lane, K., Van Valen, D., DeFelice, M.M., Macklin, D.N., Kudo, T., Jaimovich, A., Carr, A., Meyer, T., Pe'er, D., Boutet, S.C., and Covert, M.W. (2017). Measuring Signaling and RNA-Seq in the Same Cell Links Gene Expression to Dynamic Patterns of NF- κ B Activation. *Cell Syst.* *4*, 458–469.
- Lane, K., Andres-Terre, M., Kudo, T., Monack, D.M., and Covert, M.W. (2019). Escalating Threat Levels of Bacterial Infection Can Be Discriminated by Distinct MAPK and NF- κ B Signaling Dynamics in Single Host Cells. *Cell Syst.* *8*, 183–196.
- Lee, R.E.C., Walker, S.R., Savery, K., Frank, D.A., and Gaudet, S. (2014). Fold change of nuclear NF- κ B determines TNF-induced transcription in single cells. *Mol. Cell* *53*, 867–879.
- Leifer, C.A., and Medvedev, A.E. (2016). Molecular mechanisms of regulation of Toll-like receptor signaling. *J. Leukoc. Biol.* *100*, 927–941.
- Levine, J.H., Lin, Y., and Elowitz, M.B. (2013). Functional roles of pulsing in genetic circuits. *Science* *342*, 1193–1200.
- Li, P., and Elowitz, M.B. (2019). Communication codes in developmental signaling pathways. *Development* *146*, dev170977.
- Lin, B., Dutta, B., and Fraser, I.D.C. (2017). Systematic Investigation of Multi-TLR Sensing Identifies Regulators of Sustained Gene Activation in Macrophages. *Cell Syst.* *5*, 25–37.
- Lin, J., Jordi, C., Son, M., Van Phan, H., Drayman, N., Abasiyanik, M.F., Vistain, L., Tu, H.-L., and Tay, S. (2019). Ultra-sensitive digital quantification of proteins and mRNA in single cells. *Nat. Commun.* *10*, 3544.
- Litvak, V., Ramsey, S.A., Rust, A.G., Zak, D.E., Kennedy, K.A., Lampano, A.E., Nykter, M., Shmulevich, I., and Aderem, A. (2009). Function of C/EBP δ in a regulatory circuit that discriminates between transient and persistent TLR4-induced signals. *Nat. Immunol.* *10*, 437–443.
- Liu, S., Chen, J., Cai, X., Wu, J., Chen, X., Wu, Y.-T., Sun, L., and Chen, Z.J. (2013). MAVS recruits multiple ubiquitin E3 ligases to activate antiviral signaling cascades. *eLife* *2*, e00785.
- Liu, D., Sheng, C., Gao, S., Yao, C., Li, J., Jiang, W., Chen, H., Wu, J., Pan, C., Chen, S., and Huang, W. (2015a). SOCS3 Drives Proteasomal Degradation of TBK1 and Negatively Regulates Antiviral Innate Immunity. *Mol. Cell. Biol.* *35*, 2400–2413.
- Liu, S., Cai, X., Wu, J., Cong, Q., Chen, X., Li, T., Du, F., Ren, J., Wu, Y.T., Grishin, N.V., and Chen, Z.J. (2015b). Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3 activation. *Science* *347*, aaa2630.
- Livnah, O., Johnson, D.L., Stura, E.A., Farrell, F.X., Barbone, F.P., You, Y., Liu, K.D., Goldsmith, M.A., He, W., Krause, C.D., et al. (1998). An antagonist peptide-EPO receptor complex suggests that receptor dimerization is not sufficient for activation. *Nat. Struct. Biol.* *5*, 993–1004.
- Long, L., Deng, Y., Yao, F., Guan, D., Feng, Y., Jiang, H., Li, X., Hu, P., Lu, X., Wang, H., et al. (2014). Recruitment of phosphatase PP2A by RACK1 adaptor protein deactivates transcription factor IRF3 and limits type I interferon signaling. *Immunity* *40*, 515–529.
- Longo, D.M., Selimkhanov, J., Kearns, J.D., Hasty, J., Hoffmann, A., and Tsimring, L.S. (2013). Dual delayed feedback provides sensitivity and robustness to the NF- κ B signaling module. *PLoS Comput. Biol.* *9*, e1003112.
- Luo, L., Lucas, R.M., Liu, L., and Stow, J.L. (2019). Signalling, sorting and scaffolding adaptors for Toll-like receptors. *J. Cell Sci.* *133*, jcs239194.
- Ma, X., Helgason, E., Phung, Q.T., Quan, C.L., Iyer, R.S., Lee, M.W., Bowman, K.K., Starovasinik, M.A., and Dueber, E.C. (2012). Molecular basis of Tank-binding kinase 1 activation by transautophosphorylation. *Proc. Natl. Acad. Sci. USA* *109*, 9378–9383.
- Mahtani, K.R., Brook, M., Dean, J.L., Sully, G., Saklatvala, J., and Clark, A.R. (2001). Mitogen-activated protein kinase p38 controls the expression and posttranslational modification of tristetraprolin, a regulator of tumor necrosis factor alpha mRNA stability. *Mol. Cell. Biol.* *21*, 6461–6469.
- Manes, N.P., Angermann, B.R., Koppenol-Raab, M., An, E., Sjoelund, V.H., Sun, J., Ishii, M., Germain, R.N., Meier-Schellersheim, M., and Nita-Lazar, A. (2015). Targeted Proteomics-Driven Computational Modeling of Macrophage S1P Chemosensing. *Mol. Cell. Proteomics* *14*, 2661–2681.
- Mann, M., Mehta, A., Zhao, J.L., Lee, K., Marinov, G.K., Garcia-Flores, Y., Lu, L.-F., Rudensky, A.Y., and Baltimore, D. (2017). An NF- κ B-microRNA regulatory network tunes macrophage inflammatory responses. *Nat. Commun.* *8*, 851.
- Mannon, N.M., Greenwood, S.M., Young, R., Cox, S., Brindle, J., Read, D., Nelläker, C., Vesely, C., Ponting, C.P., McLaughlin, P.J., et al. (2014). The RNA-editing enzyme ADAR1 controls innate immune responses to RNA. *Cell Rep.* *9*, 1482–1494.
- Martins, A.J., Narayanan, M., Prüstel, T., Fixsen, B., Park, K., Gottschalk, R.A., Lu, Y., Andrews-Pfannkoch, C., Lau, W.W., Wendelsdorf, K.V., and Tsang, J.S. (2017). Environment Tunes Propagation of Cell-to-Cell Variation in the Human Macrophage Gene Network. *Cell Syst.* *4*, 379–392.
- McGettrick, A.F., and O'Neill, L.A.J. (2010). Localisation and trafficking of Toll-like receptors: an important mode of regulation. *Curr. Opin. Immunol.* *22*, 20–27.
- Meier-Schellersheim, M., Xu, X., Angermann, B., Kunkel, E.J., Jin, T., and Germain, R.N. (2006). Key role of local regulation in chemosensing revealed by a new molecular interaction-based modeling method. *PLoS Comput. Biol.* *2*, e82.
- Merika, M., Williams, A.J., Chen, G., Collins, T., and Thanos, D. (1998). Recruitment of CBP/p300 by the IFN β enhanceosome is required for synergistic activation of transcription. *Mol. Cell* *1*, 277–287.
- Mikuda, N., Kolesnichenko, M., Beaudette, P., Popp, O., Uyar, B., Sun, W., Tufan, A.B., Perder, B., Akalin, A., Chen, W., et al. (2018). The I κ B kinase complex is a regulator of mRNA stability. *EMBO J.* *37*, e98658.
- Mitchell, S., Vargas, J., and Hoffmann, A. (2016). Signaling via the NF κ B system. *Wiley Interdiscip. Rev. Syst. Biol. Med.* *8*, 227–241.
- Monie, T.P., Moncrieffe, M.C., and Gay, N.J. (2009). Structure and regulation of cytoplasmic adapter proteins involved in innate immune signaling. *Immunol. Rev.* *227*, 161–175.
- Morris, M.K., Saez-Rodriguez, J., Sorger, P.K., and Lauffenburger, D.A. (2010). Logic-based models for the analysis of cell signaling networks. *Biochemistry* *49*, 3216–3224.
- Natoli, G. (2009). Control of NF-kappaB-dependent transcriptional responses by chromatin organization. *Cold Spring Harb. Perspect. Biol.* *1*, a000224.
- Nau, G.J., Richmond, J.F.L., Schlesinger, A., Jennings, E.G., Lander, E.S., and Young, R.A. (2002). Human macrophage activation programs induced by bacterial pathogens. *Proc. Natl. Acad. Sci. USA* *99*, 1503–1508.
- Nelson, D.E., Ihekweba, A.E., Elliott, M., Johnson, J.R., Gibney, C.A., Foreman, B.E., Nelson, G., See, V., Horton, C.A., Spiller, D.G., et al. (2004). Oscillations in NF-kappaB signaling control the dynamics of gene expression. *Science* *306*, 704–708.
- Netea, M.G., Joosten, L.A.B., Latz, E., Mills, K.H.G., Natoli, G., Stunnenberg, H.G., O'Neill, L.A., and Xavier, R.J. (2016). Trained immunity: A program of innate immune memory in health and disease. *Science* *352*, aaf1098, aaf1098.
- Newton, K., and Dixit, V.M. (2012). Signaling in innate immunity and inflammation. *Cold Spring Harb. Perspect. Biol.* *4*, a006049, a006049.
- Ni, G., Konno, H., and Barber, G.N. (2017). Ubiquitination of STING at lysine 224 controls IRF3 activation. *Sci. Immunol.* *2*, eaah7119.

- Ni, G., Ma, Z., Wong, J.P., Zhang, Z., Cousins, E., Major, M.B., and Damania, B. (2020). PPP6C Negatively Regulates STING-Dependent Innate Immune Responses. *MBio* *11*, e01728-20.
- Nolen, B., Taylor, S., and Ghosh, G. (2004). Regulation of protein kinases; controlling activity through activation segment conformation. *Mol. Cell* *15*, 661–675.
- Ohto, U., and Shimizu, T. (2016). Structural aspects of nucleic acid-sensing Toll-like receptors. *Biophys. Rev.* *8*, 33–43.
- Ourthiague, D.R., Birnbaum, H., Ortenl f, N., Vargas, J.D., Wollman, R., and Hoffmann, A. (2015). Limited specificity of IRF3 and ISGF3 in the transcriptional innate-immune response to double-stranded RNA. *J. Leukoc. Biol.* *98*, 119–128.
- Pandey, S., Gruenbaum, A., Kanashova, T., Mertins, P., Cluzel, P., and Chevrier, N. (2020). Pairwise Stimulations of Pathogen-Sensing Pathways Predict Immune Responses to Multi-adjuvant Combinations. *Cell Syst.* *11*, 495–508.
- Panne, D., Maniatis, T., and Harrison, S.C. (2007). An atomic model of the interferon- β enhanceosome. *Cell* *129*, 1111–1123.
- Park, B.S., and Lee, J.-O. (2013). Recognition of lipopolysaccharide pattern by TLR4 complexes. *Exp. Mol. Med.* *45*, e66, e66.
- Park, S.H., Kang, K., Giannopoulou, E., Qiao, Y., Kang, K., Kim, G., Park-Min, K.-H., and Ivashkiv, L.B. (2017). Type I interferons and the cytokine TNF cooperatively reprogram the macrophage epigenome to promote inflammatory activation. *Nat. Immunol.* *18*, 1104–1116.
- Parvatiyar, K., Pindado, J., Dev, A., Aliyari, S.R., Zaver, S.A., Gerami, H., Chapon, M., Ghaffari, A.A., Dhingra, A., and Cheng, G. (2018). A TRAF3-NIK module differentially regulates DNA vs RNA pathways in innate immune signaling. *Nat. Commun.* *9*, 2770.
- Pattison, M.J., Mitchell, O., Flynn, H.R., Chen, C.-S., Yang, H.-T., Ben-Addi, H., Boeing, S., Snijders, A.P., and Ley, S.C. (2016). TLR and TNF-R1 activation of the MKK3/MKK6-p38 α axis in macrophages is mediated by TPL-2 kinase. *Biochem. J.* *473*, 2845–2861.
- Pauls, E., Shpiro, N., Peggie, M., Young, E.R., Sorcek, R.J., Tan, L., Choi, H.G., and Cohen, P. (2012). Essential role for IKK β in production of type 1 interferons by plasmacytoid dendritic cells. *J. Biol. Chem.* *287*, 19216–19228.
- Polley, S., Huang, D.-B., Hauenstein, A.V., Fusco, A.J., Zhong, X., Vu, D., Schr felfbauer, B., Kim, Y., Hoffmann, A., Verma, I.M., et al. (2013). A structural basis for I κ B kinase 2 activation via oligomerization-dependent trans autophosphorylation. *PLoS Biol.* *11*, e1001581.
- Purvis, J.E., and Lahav, G. (2013). Encoding and decoding cellular information through signaling dynamics. *Cell* *152*, 945–956.
- Rahighi, S., Ikeda, F., Kawasaki, M., Akutsu, M., Suzuki, N., Kato, R., Kensche, T., Uejima, T., Bloor, S., Komander, D., et al. (2009). Specific recognition of linear ubiquitin chains by NEMO is important for NF- κ B activation. *Cell* *136*, 1098–1109.
- Ramirez-Carrozzi, V.R., Braas, D., Bhatt, D.M., Cheng, C.S., Hong, C., Doty, K.R., Black, J.C., Hoffmann, A., Carey, M., and Smale, S.T. (2009). A unifying model for the selective regulation of inducible transcription by CpG islands and nucleosome remodeling. *Cell* *138*, 114–128.
- Roberts, A.W., Lee, B.L., Deguine, J., John, S., Shlomchik, M.J., and Barton, G.M. (2017). Tissue-Resident Macrophages Are Locally Programmed for Silent Clearance of Apoptotic Cells. *Immunity* *47*, 913–927.
- Roget, K., Ben-Addi, A., Mambole-Dema, A., Gantke, T., Yang, H.-T., Janzen, J., Morrice, N., Abbott, D., and Ley, S.C. (2012). I κ B kinase 2 regulates TPL-2 activation of extracellular signal-regulated kinases 1 and 2 by direct phosphorylation of TPL-2 serine 400. *Mol. Cell. Biol.* *32*, 4684–4690.
- Rothchild, A.C., Sissons, J.R., Shafiani, S., Plaisier, C., Min, D., Mai, D., Gilchrist, M., Peschon, J., Larson, R.P., Bergthaler, A., et al. (2016). MiR-155-regulated molecular network orchestrates cell fate in the innate and adaptive immune response to *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA* *113*, E6172–E6181.
- Saccani, S., Pantano, S., and Natoli, G. (2001). Two waves of nuclear factor κ B recruitment to target promoters. *J. Exp. Med.* *193*, 1351–1359.
- Salojin, K.V., Owusu, I.B., Millerchip, K.A., Potter, M., Platt, K.A., and Oravec, T. (2006). Essential role of MAPK phosphatase-1 in the negative control of innate immune responses. *J. Immunol.* *176*, 1899–1907.
- Sasai, M., Linehan, M.M., and Iwasaki, A. (2010). Bifurcation of Toll-like receptor 9 signaling by adaptor protein 3. *Science* *329*, 1530–1534.
- Satija, R., and Shalek, A.K. (2014). Heterogeneity in immune responses: from populations to single cells. *Trends Immunol.* *35*, 219–229.
- Scholes, C., DePace, A.H., and S nchez,  . (2017). Combinatorial Gene Regulation through Kinetic Control of the Transcription Cycle. *Cell Syst.* *4*, 97–108.
- Scholz, R., Sidler, C.L., Thali, R.F., Winssinger, N., Cheung, P.C.F., and Neumann, D. (2010). Autoactivation of transforming growth factor β -activated kinase 1 is a sequential bimolecular process. *J. Biol. Chem.* *285*, 25753–25766.
- Schr felfbauer, B., and Hoffmann, A. (2011). How do pleiotropic kinase hubs mediate specific signaling by TNFR superfamily members? *Immunol. Rev.* *244*, 29–43.
- Schr felfbauer, B., Polley, S., Behar, M., Ghosh, G., and Hoffmann, A. (2012). NEMO ensures signaling specificity of the pleiotropic IKK β by directing its kinase activity toward I κ B α . *Mol. Cell* *47*, 111–121.
- Sen, S., Cheng, Z., Sheu, K.M., Chen, Y.H., and Hoffmann, A. (2020). Gene Regulatory Strategies that Decode the Duration of NF κ B Dynamics Contribute to LPS- versus TNF-Specific Gene Expression. *Cell Syst.* *10*, 169–182.
- Sheu, K., Luecke, S., and Hoffmann, A. (2019). Stimulus-specificity in the responses of immune sentinel cells. *Curr. Opin. Syst. Biol.* *78*, 53–61.
- Shih, V.F.-S., Kearns, J.D., Basak, S., Savinova, O.V., Ghosh, G., and Hoffmann, A. (2009). Kinetic control of negative feedback regulators of NF- κ B/RelA determines their pathogen- and cytokine-receptor signaling specificity. *Proc. Natl. Acad. Sci. USA* *106*, 9619–9624.
- Shim, J.-H., Xiao, C., Paschal, A.E., Bailey, S.T., Rao, P., Hayden, M.S., Lee, K.-Y., Bussey, C., Steckel, M., Tanaka, N., et al. (2005). TAK1, but not TAB1 or TAB2, plays an essential role in multiple signaling pathways in vivo. *Genes Dev.* *19*, 2668–2681.
- Shinohara, H., Behar, M., Inoue, K., Hiroshima, M., Yasuda, T., Nagashima, T., Kimura, S., Sanjo, H., Maeda, S., Yumoto, N., et al. (2014). Positive feedback within a kinase signaling complex functions as a switch mechanism for NF- κ B activation. *Science* *344*, 760–764.
- Shu, C., Sankaran, B., Chaton, C.T., Herr, A.B., Mishra, A., Peng, J., and Li, P. (2013). Structural insights into the functions of TBK1 in innate antimicrobial immunity. *Structure* *21*, 1137–1148.
- Skaug, B., Chen, J., Du, F., He, J., Ma, A., and Chen, Z.J. (2011). Direct, non-catalytic mechanism of IKK inhibition by A20. *Mol. Cell* *44*, 559–571.
- Sparrer, K.M., and Gack, M.U. (2015). Intracellular detection of viral nucleic acids. *Curr. Opin. Microbiol.* *26*, 1–9.
- Spencer, S.L., Gaudet, S., Albeck, J.G., Burke, J.M., and Sorger, P.K. (2009). Non-genetic origins of cell-to-cell variability in TRAIL-induced apoptosis. *Nature* *459*, 428–432.
- Stempel, M., Chan, B., Jurani  Lisni , V., Krmpotić, A., Hartung, J., Paludan, S.R., F llbrunn, N., Lemmermann, N.A., and Brinkmann, M.M. (2019). The herpesviral antagonist m152 reveals differential activation of STING-dependent IRF and NF- κ B signaling and STING’s dual role during MCMV infection. *EMBO J.* *38*, e100983.
- Sung, M.-H., Li, N., Lao, Q., Gottschalk, R.A., Hager, G.L., and Fraser, I.D.C. (2014). Switching of the relative dominance between feedback mechanisms in lipopolysaccharide-induced NF- κ B signaling. *Sci. Signal.* *7*, ra6, ra6.
- Takeuchi, O., and Akira, S. (2010). Pattern recognition receptors and inflammation. *Cell* *140*, 805–820.
- Tan, R.S.T., Ho, B., Leung, B.P., and Ding, J.L. (2014). TLR cross-talk confers specificity to innate immunity. *Int. Rev. Immunol.* *33*, 443–453.
- Tay, S., Hughey, J.J., Lee, T.K., Lipniacki, T., Quake, S.R., and Covert, M.W. (2010). Single-cell NF- κ B dynamics reveal digital activation and analogue information processing. *Nature* *466*, 267–271.

- Testa, U., Pelosi, E., Castelli, G., and Labbaye, C. (2017). miR-146 and miR-155: Two Key Modulators of Immune Response and Tumor Development. *Noncoding RNA* 3, 22.
- Tokunaga, F. (2013). Linear ubiquitination-mediated NF- κ B regulation and its related disorders. *J. Biochem.* 154, 313–323.
- Tomida, T., Takekawa, M., and Saito, H. (2015). Oscillation of p38 activity controls efficient pro-inflammatory gene expression. *Nat. Commun.* 6, 8350.
- Tong, A.J., Liu, X., Thomas, B.J., Lissner, M.M., Baker, M.R., Senagolage, M.D., Allred, A.L., Barish, G.D., and Smale, S.T. (2016). A Stringent Systems Approach Uncovers Gene-Specific Mechanisms Regulating Inflammation. *Cell* 165, 165–179.
- Turner, D.A., Paszek, P., Woodcock, D.J., Nelson, D.E., Horton, C.A., Wang, Y., Spiller, D.G., Rand, D.A., White, M.R.H., and Harper, C.V. (2010). Physiological levels of TNF α stimulation induce stochastic dynamics of NF- κ B responses in single living cells. *J. Cell Sci.* 123, 2834–2843.
- Uematsu, S., Sato, S., Yamamoto, M., Hirohata, T., Kato, H., Takeshita, F., Matsuda, M., Coban, C., Ishii, K.J., Kawai, T., et al. (2005). Interleukin-1 receptor-associated kinase-1 plays an essential role for Toll-like receptor (TLR)7- and TLR9-mediated interferon- α induction. *J. Exp. Med.* 201, 915–923.
- Vanaja, K.G., Timp, W., Feinberg, A.P., and Levchenko, A. (2018). A Loss of Epigenetic Control Can Promote Cell Death through Reversing the Balance of Pathways in a Signaling Network. *Mol. Cell* 72, 60–70.
- Voisinne, G., Nixon, B.G., Melbinger, A., Gasteiger, G., Vergassola, M., and Altan-Bonnet, G. (2015). T Cells Integrate Local and Global Cues to Discriminate between Structurally Similar Antigens. *Cell Rep.* 11, 1208–1219.
- Wang, C., Deng, L., Hong, M., Akkaraju, G.R., Inoue, J., and Chen, Z.J. (2001). TAK1 is a ubiquitin-dependent kinase of MKK and IKK. *Nature* 412, 346–351.
- Waterfield, M.R., Zhang, M., Norman, L.P., and Sun, S.C. (2003). NF- κ B1/p105 regulates lipopolysaccharide-stimulated MAP kinase signaling by governing the stability and function of the Tpl2 kinase. *Mol. Cell* 11, 685–694.
- Werner, S.L., Barken, D., and Hoffmann, A. (2005). Stimulus specificity of gene expression programs determined by temporal control of IKK activity. *Science* 309, 1857–1861.
- Werner, S.L., Kearns, J.D., Zadorozhnaya, V., Lynch, C., O’Dea, E., Boldin, M.P., Ma, A., Baltimore, D., and Hoffmann, A. (2008). Encoding NF- κ B temporal control in response to TNF: distinct roles for the negative regulators I κ B α and A20. *Genes Dev.* 22, 2093–2101.
- Wertz, I.E., O’Rourke, K.M., Zhou, H., Eby, M., Aravind, L., Seshagiri, S., Wu, P., Wiesmann, C., Baker, R., Boone, D.L., et al. (2004). De-ubiquitination and ubiquitin ligase domains of A20 downregulate NF- κ B signalling. *Nature* 430, 694–699.
- Witzel, F., Maddison, L., and Blüthgen, N. (2012). How scaffolds shape MAPK signaling: what we know and opportunities for systems approaches. *Front. Physiol.* 3, 475.
- Wu, J., Sun, L., Chen, X., Du, F., Shi, H., Chen, C., and Chen, Z.J. (2013). Cyclic GMP-AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* 339, 826–830.
- Xia, Z.-P., Sun, L., Chen, X., Pineda, G., Jiang, X., Adhikari, A., Zeng, W., and Chen, Z.J. (2009). Direct activation of protein kinases by unanchored polyubiquitin chains. *Nature* 461, 114–119.
- Xia, C., Fan, J., Emanuel, G., Hao, J., and Zhuang, X. (2019). Spatial transcriptome profiling by MERFISH reveals subcellular RNA compartmentalization and cell cycle-dependent gene expression. *Proc. Natl. Acad. Sci. USA* 116, 19490–19499.
- Xiang, W., Zhang, Q., Lin, X., Wu, S., Zhou, Y., Meng, F., Fan, Y., Shen, T., Xiao, M., Xia, Z., et al. (2016). PPM1A silences cytosolic RNA sensing and antiviral defense through direct dephosphorylation of MAVS and TBK1. *Sci. Adv.* 2, e1501889.
- Xu, X., Meckel, T., Brzostowski, J.A., Yan, J., Meier-Schellersheim, M., and Jin, T. (2010). Coupling mechanism of a GPCR and a heterotrimeric G protein during chemoattractant gradient sensing in Dictyostelium. *Sci. Signal.* 3, ra71, ra71.
- Yoboua, F., Martel, A., Duval, A., Mukawera, E., and Grandvaux, N. (2010). Respiratory syncytial virus-mediated NF- κ B p65 phosphorylation at serine 536 is dependent on RIG-I, TRAF6, and IKK β . *J. Virol.* 84, 7267–7277.
- Zanoni, I., and Granucci, F. (2012). Regulation and dysregulation of innate immunity by NFAT signaling downstream of pattern recognition receptors (PRRs). *Eur. J. Immunol.* 42, 1924–1931.
- Zanoni, I., Ostuni, R., Marek, L.R., Barresi, S., Barbalat, R., Barton, G.M., Granucci, F., and Kagan, J.C. (2011). CD14 controls the LPS-induced endocytosis of Toll-like receptor 4. *Cell* 147, 868–880.
- Zhang, J., Clark, K., Lawrence, T., Pegg, M.W., and Cohen, P. (2014). An unexpected twist to the activation of IKK β : TAK1 primes IKK β for activation by autophosphorylation. *Biochem. J.* 461, 531–537.
- Zhang, L., Wei, N., Cui, Y., Hong, Z., Liu, X., Wang, Q., Li, S., Liu, H., Yu, H., Cai, Y., et al. (2018). The deubiquitinase CYLD is a specific checkpoint of the STING antiviral signaling pathway. *PLoS Pathog.* 14, e1007435.
- Zhang, C., Tu, H.-L., Jia, G., Mukhtar, T., Taylor, V., Rzhetsky, A., and Tay, S. (2019a). Ultra-multiplexed analysis of single-cell dynamics reveals logic rules in differentiation. *Sci. Adv.* 5, eaav7959.
- Zhang, C., Shang, G., Gui, X., Zhang, X., Bai, X.-C., and Chen, Z.J. (2019b). Structural basis of STING binding with and phosphorylation by TBK1. *Nature* 567, 394–398.
- Zhao, Y., Liang, L., Fan, Y., Sun, S., An, L., Shi, Z., Cheng, J., Jia, W., Sun, W., Mori-Akiyama, Y., et al. (2012). PPM1B negatively regulates antiviral response via dephosphorylating TBK1. *Cell. Signal.* 24, 2197–2204.